Traces of DNA evidence can be magnified with PCR
DNA typing
Identification of unknown substances for the presence of drugs with color indicator tests
Urinalysis for drugs by thin layer chromatography

Nichole Kellerman
Nichole_A_Kellerman@mcpsmd.org
Walter Johnson High School
6400 Rock Spring Drive
Bethesda, MD 20814

Mentor
Pamela Schwartzberg, M.D., Ph.D.
National Human Genome Research Institute, National Institute of Health

I. Maryland State Standards :

1.1.1 The student will recognize that real problems have more than one solution and decisions to accept one solution over another are made on the basis of many issues.
1.1.2 The student will modify or affirm scientific ideas according to accumulated evidence.
1.2.2 The student will identify meaningful, answerable, scientific questions.
1.2.3 The student will pose meaningful, answerable, scientific questions.
1.2.6 The student will identify appropriate methods for conducting an investigation (independent and dependent variables, proper controls, repeat trials, appropriate sample size etc.)
1.3.1 The student will develop and demonstrate skills in using lab and field equipment to perform investigative techniques.
1.3.2 The student will recognize safe laboratory procedures.
1.3.3 The student will demonstrate safe handling of the chemicals and materials of science.
1.3.4 The student will learn the use of new instruments and equipment by following instructions in a manual or fro

II. Unit Goals:

Students will become aware of immunology concepts used to create the technology Forensic Technicians use in crime labs. Students will investigate a crimescene, examine trace evidence for DNA, identify unknown white powders, and examine urine samples from possible suspects. They will use DNA extraction to obtain DNA from the hair follicle and PCR to magnify the suspects' DNA samples. Students will determine the specific analytic test that should be performed on each item of evidence collected, in order to identify which suspect committed the crime. Choices of evaluation tools will be various tests for drugs or pregnancy, and gel electrophoresis.

After completing this unit, students will be able

IV. Science Background and Teacher Resources:

A PowerP

dextromethorphan/ pseudoephedrine, Dexatrim, Sudafed, or Quinine. The "control" and unknown urine samples are added to a Tox Elut® column filled with diatomaceous Earth to ex

V. Learning Objectives

1x TBE buffer	1000 mL	This should be enough for 3 classes to make gels and to use as gel running bugger. This is a 10% Solution.
P20 & P200 uL micropipettes	8 each	
1.5 mL microcentrifuge tubes	16 per class	
Timer	1	Students can use timers on their phones. Teacher may also set a timer on the board for all students to use.

MicropiptC q 40.5(ic)-6001 Tc 0

Lessons 6 & 7 Materials	Quantity	Comments
Aspirin	1 box	Use a mortar and pestle to ground the entire box into a powder,

This is what each class set should look like.

Labels for dropper bottles

Methanol	Methanol	Methanol	Methanol	Methanol	Methanol	Methanol	Methanol
Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M	Hydrochloric Acid 3.0M
Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M	Iron (III) Nitrate 0.1M
Universal Indicator	Universal Indicator	Universal Indicator	Universal Indicator	Universal Indicator	Universal Indicator	Universal Indicator	Universal Indicator

CONTACT DEXTROMETHORPHAN/ PSEUDOEPHEDRINE	CAFFEINE	DEXATRIM PHENYLPROPANOLAMI	SUDAFED PSEUDOEPHEDRI	QUININE	COMBINED STANDARD	COMBINED CONTROL
UNKNOWN #1	UNKNOWN # 2	UNKNOWN #3	UNKNOWN # 4	UNKNOWN #5	UNKNOWN # 6	

Lessons 8 & 9 Materials	Quantity	Comments
Methanol		Use methanol to add to water to create a 50:50 mixture to dissolve each chemical listed.
Ring stands	6	Attach test tube clamps to ring stands
Test tube ring stand clamps	6	

Caffeine
(caffedrn B0(nd)-ET EMC /P <</

Sudafed = Pseudoephedrine	1 open capsule	Add one capsule to100 mL of distilled water and 100mL of methanol.
Quinine	1 open capsule	Add one capsule to100 mlof distilled water and 100mL of methanol.
Dexatrim	1 open capsule	Add one capsule to100 mL of distilled water and 100mL of methanol.
Urine or synthetic urine can be used	150 mL	Needed per class. You can use urine or purchase synthetic urine.
7 glassbottles with caps	7	
6 ring stands	6	
6 silica gel plates	6	
6 test tube holder clamps to go onto the ring stand	6	
6 capillary tubes	6	
6 tox elute column Filled with diatomaceous Earth	6	
6 spray bottles	6	
6 50 ml beakers	6	
Hair dryer	1	
Eluent solution	2 bottles	
Ninhydrin solution	1 bottle	
Mercury oxide solution		
Rf value cards	6	
Isopropanol		
Simulated urine	1 bottles	1 bottle per class
0,1 N HCl		
Pipettes	6	
Water bath		
Hood		
UV lamps black light	1 or 7	Groups can share one. There are little UV flashlights that you can give each students
Acetone		
Ethyl acetate		
Ammonium hydroxide		
Methanol		
Chloroplantinic acid		
Sulfuric acid 10% solution		
Ethyl acetate		
25mL Graduated cylinders	6	
16 ozglass Mason jar with a lid	6 or 7	The chromatography plateis inside of the Mason jar
Capillary tubes	6 per class	
Blow dryer	1	
Combined standard		NO URINE!!! 10mL of each of these

VIII. Teacher Guide

Lesson 1

- Show students the Power Point slides # 1-23.
-

Lesson 3:

- Students will put on gloves, goggles, and aprons .
- Students will receive a tray containing all the supplies for DNA extraction.
- Students will use tweezers to remove hair from two members. Each microcentrifuge tube used for the DNA extraction will receive a number # 1-16 so that students will not be able to know the names of each DNA extraction.

I. Isolation of Genomic DNA from Hair

1. Obtain 3 to 4 hairs with root and sheath attached. Cut shafts of hair to approximately 2 cm lengths and put in 1.5 ml microcentrifuge tube.
2. Shake Chelex solution. While beads are still suspended, remove 150 uL and add to hairs in microcentrifuge tube. Cellular components contain heavy metal ions can interfere with Taq polymerase in PCR reaction. Chelex binds these metal ions and any DNAases that will degrade genomic DNA.
3. Place mixtures in vortex or swirl vigorously. Make sure hairs are in the solution.
4. Incubate at 56 °C for 5 minutes
5. Place mixtures in vortex or swirl vigorously
   Make sure hairs are in the solution.
6. Incubate at 56° C for another 10 minutes.
7. Place mixtures in vortex or swirl vigorously. Make sure hairs are still re-suspended.
8. Incubate in boiling water for 10 minutes. This breaks the cell open.
9. Place mixtures in vortex or swirl vigorously and then put on ice for 5 minutes.
10. Spin in microcentrifuge for 5 minutes
11. Slowly and carefully remove tube. DO NOT disturb the pellet at bottom of tube.
12. Remove 75 uL of supernatant and place in a fresh tube. DO NOT transfer any Chelex beads.
13. Label tubes and place on ice or store in freezer until ready to continue.

Make sure the PCR machine has been programmed.

Thermocycler settings: 30 cycles

- 72 C for 2 minutes: Taq polymerase will add new bases to make the complementary strands

II. PCR Reaction Set Up and Amplification

1. Use a permanent marker to label a PCR tube supplied by your teacher. It will contain a white bead that contains reagents for the amplification reaction. Make sure the bead dissolves when the DNA is added to the tube
2. Use the matrix below as a checklist while added reagents to the PCR tube.

Write on the microcentrifuge tube	Hair Cell DNA	Upstream primer	Downstream primer	Total volume
Period #____ 3 Initials of your name	20 uL	2.5 uL	2.5 uL	25 uL

3. Close tube cap tightly and place in block of automated thermal cycler.

NOTE: Keep previous class sets in the refrigerator or on ice until all your classes are finished. You can load all classes into the Thermocycler at the same time and then you can leave them there all night.

Lesson 4:

- Students will put on gloves, goggles, and n Pek samp(t)-le udec4(r o. Sa)-4(r)2(e)2(-2(e)2(n)5

5. Is this individual or class evidence? Explain your answer.
   Examples of individual evidence are fingerprints, handwriting, DNA patterns, and sometimes physical matches, However, most evidence is class evidence, characteristics common to a group of similar objects, not to one single object.
6. Comment on the Frye Standard; the "general acceptance" of this scientific technique. Would your scientific investigation hold up in a court of law if you were asked to take the stand? Please include a discussion of your training, the number of times you have done this technique, your sterile technique (food, gloves, sterilized lab bench), and laboratory procedures you followed (any possible mistakes made during procedure, identify possible steps in procedure that you may have accidentally contaminated your sample)?
   Frye Standard Summary: Defendant James Alphonso Frye was convicted of murder in the second degree and appealed the decision. The issue presented for consideration is that defense counsel offered an expert witness to testify to the result of a systolic blood pressure deception test, a rudimentary precursor to the lie detector, and was denied. They further offered that a test be conducted in the courtroom and were again denied. The prosecution argued that "while the courts will go a long way in admitting expert testimony, deduced from a well recognized scientific principle or discovery, the thing from which the deduction is made must be

Lesson 6-7:

- Show students PowerPoint slides #37- 43.
- All students should put on gloves, goggles, and laboratory aprons.
- Use a mortar and pestle to ground up these pills:
  - Aspirin
  - Baking soda (sodium bick 0 Td [(Pick)6(0 Td [(Pick)6(0 Td [(Pick)6(0 Td [(Pig)6(late Tw

Lesson 8:

Teacher preparation before the lesson:

- Show students PowerPoint slides #44 – 54.
- Use a test tube clamp mounted on a ring stand to support each column used. Place a collection beaker under each column. A 50 or 100 mL beaker should work. A maximum of 50 mL of Eluent solution is collected.
- Make 5 different individual "standard" solutions.
    - Dilute the drug samples with 100mL water 100mL methanol mixture. Place each in a glass bottle.
        - Sudafed
        - Contact
        - Caffeine pills (or Caffedrine)
        - Dexatrim
        - Quinine
        - Add 20 mL of urine sample and 50 mL of each drug solution.
        - 6 unknowns need to be made. Place one drug in each sample. You can mix 2 drugs in 2 of the unknowns or leave one unknown without drugs.
        - Label these mixtures unknown # 1-6
- Make a control sample that has 60mL of urine without any drugs
- Make a combined standard by placing 10 mL of each of the 5 samples. Add 20 mL of urine to this combined standard.
- Place the water bath inside the fume hood. The vapors released from these samples are toxic and should only be handled within a fume hood.

During the lesson:

- Have students read the introduction to the lab
- Teacher will pour 20 mL of the control into the apparatus.
- Have students pour 20mL of their unknown into their apparatus.
- Show students PowerPoint slides #55 – 57.

10. Place the collection beakers in a well ventilated area and allow them to evaporate overnight.
    If this is not possible because there is not enough time, place the collection beakers under a hood into the water bath at approximately 65-70°C. Allow the beakers to evaporate until they are dry.

Lesson 9:

Part II: Prepare the developing chamber and the chromatography plate.

1. Take a 16 oz glass Mason jar with a lid. The jar will hold the 2" x 4" chromatography plate.
2. The Teacher will prepare the developing solvent by adding ethyl acetate, methanol, and concentrated ammonium hydroxide together in a ratio of 17:2:1.
    - Get a small bottle or a 50-100 mL beaker.
    - Add 34 mL ethyl acetate.
    - Add 4 mL methanol
    - Add 2 mL concentrated ammonium hydroxide.
    - This will be sufficient for 5 mL of the developing solvent for 1 class.
    - Double these measurements and there will be 3 mL for each group of 6 set-ups. This will be enough for 3 classes
3. Add 3-5 mL of the developing solvent to each chromatography chamber. The solvent must cover the bottom of the 16 oz Mason jar Chromatography chamber.
4. Take a 5 x 10 cm chromatography plate. Use a sharp pencil to scribe 2 lines about 1.5 cm apart and down the entire 10cm length of the plate.
5. The 3 tracks will provide room for:
    - One student urine sample
    - One standard
    - One control

    The tracts keep the drug spots separated during development.

Unknown sample
Standard
Control

6. Use the pencil to make a light mark at the 9 cm height on the chromatography plate.
7. Heat the chromatography plate for 3 minutes with a blow dryer on the highest setting (90 °C or 200°C).

Part III: Reconstitution

1. Add 3 drops of methanol in each totally evaporated collection beaker to reconstitute the drug mixture.
2. Swirl the collection beaker slightly to allow the methanol to contact the entire drug residue.

Spot the unknown section of the chromatography plate:

Unknown sample
Standard
Control

3. Place the top of a 20 uL capillary tube into the methanol/drug mixture. Allow the fluid to rise as high as possible in the capillary by normal capillary attraction. This should be at least 1" in height in the tube
    a. Bring the tip of the capillary tube into the same spot on the chromatography plate about 1.5-2 cm from the bottom of the plate.
    b. Make small spots the size of a ".0
    c. Allow each spot pressed by the capillary tube to dry before allowing the capillary to redeposit its next spot on the top of the previous spot.

Spot the combined standard section of the chromatography plate:

Unknown sample
Standard
Control

4. Place the top of a 20 uL capillary into the "combined standard" mixture. Allow the fluid to rise as high as possible in the capillary by normal capillary attraction. This should be at least 1" in height in the tube .
    a. Bring the tip of the capillary tube into the same spot on the chromatography plate about 1.5-2 cm from the bottom of the plate.
    b. Make small spots the size of a ".0
    c. Allow each spot pressed by the capillary tube to dry before allowing the capillary to redeposit its next spot on the top of the previous spot.

Spot the control section of the chromatography plate:

Unknown sample
Standard
Control

5. Place the top of a 20 uL capillary into the "control" beaker. Allow the fluid to rise as high as possible in the capillary by normal capillary attraction. This should be at least 1" in height in the tube .
    a. Bring the tip of the capillary tube into the same spot on the chromatography plate about 1.5-2 cm from the bottom of the plate.
    b. Make small spots the size of a ".0
    c. Allow each spot pressed by the capillary tube to dry before allowing the capillary to redeposit its next spot on the top of the previous spot.

Now, the test chromatography plate contains samples on all three sections.

Unknown sample
Standard

Part V: Visualization

Solution	RF value	Color	Place on the chromatography plate
Ninhydrin , phenylpropanolamine	0.55		
pseudo ephedrine	0.35	Bright Pink or magenta	Brown spot
Mercury sulfate	N/A	Will turn purple before drying	
Dextromethorphan	0.55	Brown spot	Brown spot
Quinine	0.63	Brown spot Under black light it fluoresces	Brown spot highest on the chromatography plate
Caffeine	0.67	Greenish white	The highest of the three drugs on the chromatography plate
Valium and Librium		Under a black light will fluoresces	

http://www.marz -kreations.com/Chemistry/Chromatography/Dyes/RF -Values.html

IX. Student Handouts

Name: _____

Instructions:

Create a list of items that should be collected as evidence.

_____
_____
_____

How would you collect, store, and document the evidence recovered?

_____
_____
_____

What questions need to be answered to solve the crime?

DNA Fingerprinting and PCR

BACKGROUND

When Crime Scene Investigators (CSI's) arrive at the scene many times the sample left behind is not a fingerprint. Most fingerprints are partial prints and are not good enough to match a potential suspect to a crimescene. Blood or hair is readily available at crime scenes, even if found in very small samples. Before CSI's can compare DNA found at the crime scene to the DNA collected from possible suspects, they usually have to use polymerase chain reactions (PCR) to amplify the trace amounts of DNA found at the crime scene.

Forensic scientists use a technique developed by molecular geneticist Ray White in 1980, known as "DNA fingerprinting." In 1986, PCR was first used during a criminal case in the United States. The case "Pennsylvania vs. Pestinikas," involved an allegation of a funeral home in Pennsylvania that was accused of switching body parts (http://www.nfstc.org/pdi/Subject01/pdi_s01_m02_01.htm).

In 1987, DNA fingerprinting was first used by the criminal justice system in England during the "Footpath Murders." The police investigating the case were thrilled to hear what Alec Jeffreys was doing with DNA profiling and wanted to solidify their case with this new technology. They asked Alec Jeffery's to examine Richard Buckland's DNA and compare it to the DNA sample behind in one of the victims clothing. The analysis led to a conclusion the police officers had not anticipated. Instead of corroborating his testimony, the DNA analysis revealed that Richard Buckland was not a match. This evidence was used to exonerate Richard Buckland who made a false confession about raping and murdering two teenage girls after hours of interrogation by Leicester police in England. Analyzing the DNA from the victims clothing also allowed the police to know a little more about their suspect. He was a male with Type A blood, and he also had a rare enzyme found in 10% of the local male population. They eventually did get the correct man, Colin Pitchfork. The wonderful thing about DNA fingerprinting is that it can be used to prove guilt and innocence (http://www.trutv.com/library/crime/criminal_mind/forensics/dna/2.html).

CODIS is a DNA database used by police and FBI to identify DNA found at the crime scenes with known suspects. This process relies on probability. The FBI has identified 13 short tandem repeats (STR) that are useful to identify or exonerate criminals. The number of repeats within an STR is referred to as an allele of the gene in that individual. The FBI determined that probability of two unrelated Caucasians having identical alleles at all 13 STRs is approximately 1 in 575 trillion (Reilly, 2001). We will analyze one STR called the Alu insertion. When it is present it makes an allele that is approximately 400 base pairs

Remember it is just as important to eliminate a possible suspect as it is to find the correct criminal.

Well A:

Image

II. PCR Reaction Set Up and Amplification

1. Use a permanent marker to label a PCR tube supplied by your teacher. It will contain a white bead that contains reagents for the amplification reaction. Make sure the bead dissolves when the DNA is added to the tube.
2. Use the matrix below as a checklist while adding reagents to the PCR tube.

Write on the microcentrifuge tube	Hair Cell DNA	Upstream primer	Downstream primer	Total volume
Period #____ 3 Initials of your name	20 uL	2.5 uL	2.5 uL	25 uL

3. Close tube cap tightly and place in block of automated thermal cycler.

Thermocycler settings: 30 cycles

- 94 °C for 1 minute: The double stranded DNA is denatured to single strands.
- 58 °C for 2 minutes: The primers attach to specific DNA sequences of single strands.
- 72 °C for 2 minutes: Taq polymerase add new bases to make the complementary strands

III. 2.0% Agarose Gel Preparation

1. Carefully pour agarose solution into the gel casting tray to fill to a depth of 5 mm. The gel should cover about one-third of the height of the comb teeth.
2. Answer questions #-8 while you are waiting for your gel to solidify.

Name _____

Results and Discussion

1. After a photograph of your gel has been taken, orient the photograph with the sample wells at the top.
2. Attach photo in space provided and label the lanes 1 through 8, noting which lanes were designated for the ladders and which ones for student samples.

3. Using the DNA ladders as a guide, view the student sample lanes and note any allele bands.

Note: it is possible that the primers may have been amplified themselves creating a primer dimmer.

Totals			

5.

10. Comment on the Frye Standard, the "general acceptance" of this scientific technique. Would your scientific investigation hold up in a court of law if you were asked to take the stand? Please include a discussion of your training, the number of times you have done this technique, your sterile technique (food, gloves, sterilized lab bench, etc.), and laboratory procedures you followed (any possible mistakes made during procedure, identify possible steps in procedure that you may have accidentally contaminated your sample)?

RESULTS:
Identify each unknown white powder found at the crimescene. Describe the – ‡ • – • – Š ƒ – yo u Ž t ž r e a c t h i s conclusion .

Unknown #1:

Unknown #2:

Unknown #3:

Unknown #4:

Unknown #5:

What type of test did you perform?

Do you feel your group used great laboratory technique? Explain why or why not?

Explain how color indicator test may be useful to CSI out in the field.

- Drug Extraction is the urine sample containing drugs. It is poured through a column of diatomaceous earth. The diatomaceous earth has been buffered to a pH = 9. The drugs are later removed by pouring a mixture of dichloromethane, isopropanol, and ethyl acetate through the column. The drugs were previously dissolved in this mixture, the urine materials are left behind in the Tox Elut® column filled with diatomaceous Earth. The drug solution emerging from the column will appear as a spot on the chromatography plates.
- Eluent

Procedure:

Part I: Extracting the drugs from unknown urine samples

1. Make sure that you have identified your unknown solution. Unknown # _____
2. Take a look at the control solution and make sure that you have:
   - x A test tube clamp
   - x Ring stand
   - x 50 or 100 mL beaker
   - x Bottle container with unknown number matching your diatomaceous earth
   - x Graduated cylinder
3. Observe the teacher pour 20 mL of the control into the container labeled control. This sample contains no drugs.

4. Why do we need a control?

   _____

   _____

5. What is the difference between a control and an unknown?

   _____

   _____

6. Pour 20 mL of the unknown into the column containing diatomaceous earth.
7. The drugs we are trying to identify are:

   x

8. Label a 50 – 100 mL collection beaker with the unknown number.
9. Place the 50-100 mL collection beaker under the column.
10. After the columns have stopped dripping (approximately 10 minutes) remove the collection beakers from underneath the columns.
11. Place the collection beakers in a well-ventilated area and allow them to evaporate over night. If this is not possible because there is not enough time, place the collection beakers under a fume hood into the water bath at approximately 65 °C-70°C. Allow the beakers to evaporate into dryness.

Part V: Visualization

Solution	RF value	Color	Place on the chromatography plate
Ninhydrin, Phenylpropanolamine	0.55		
Pseudo ephedrine	0.35	Bright Pink or magenta	Brown spot
Mercury sulfate	N/A	Will turn purple before drying	
Dextromethorphan	0.55	Brown spot	Brown spot
Quinine	0.63	Brown spot Under black light it fluoresces	Brown spot highest on the chromatography plate
Caffeine	0.67	Greenish white	The highest of the three drugs on the chromatography plate
Valium and Librium		Under a black light will fluoresces	

Teacher will place these solutions into spray bottles, because they are toxic. They are corrosive to the skin, to the eyes, and mucous membranes.

Ninhydrin,
Iodoplatinate
Diphenylcarbasone

Mercuric sulfate is a poison .

b. Calculate the distance of a spot from the origin. Label this value SD
c. Calculate the distance of the solvent front from the origin. Label this value SF.
d. SD /SF = RF value
e. Carefully, compare the Rf values and colors of the "control" to the Rf values and colors listed on the Rf card
f. Identify all spots on the chromatography chard. If any are missing, ask your teacher to help you.

3. Identify the drugs present in the "unknown" that you have measured.
4. Share your results with the class.
5. Label the spots of all of the unknowns on the sample below.
6. Write your calculated Rf on the chart for the class.

Results

Unknown sample
Standard
Control

Sample	SF	SD	RF = SD/SF	Identification of drugs in urine
Unknown #1				
Unknown #2				

X. References

Cannons J.L., Yu L.J., Jankovic D., Crotty S., Horai R., Kirby M., Anderson S., Chevez-Barrios P., Sher A., Schwartzberg P.L. SAP regulates T cell-mediated help for humoral immunity by mechanism distinct from cytokine regulation. J. Exp. Med. 203: 1551-65 (2006). http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2118305/

Houck M.M., Siegel J.A. (2006) Fundamentals of Forensic Science 1st Ed

Yin L., Al-Alem U., Liang J., Tong W.M., Li C., Badiali M., Médard J.J., Sumegi J., Wang Z.Q., Romeo G. Mice deficient in the X-linked lymphoproliferative disease gene sap exhibit increased susceptibility to murine gammaherpesvirus-68 and hypo-gammaglobulinemia. J. Med. Virol. 71: 446-55 (2003). http://www.ncbi.nlm.nih.gov/pubmed/12966553

http://www.accessexcellence.org/AE/AEPC/DNA/detection.php

https://www.amherst.edu/system/files/media/1898/2012_Manual_5c_Alu_Lab.pdf

http://biotech.law.lsu.edu/map/TheFryeRule.html

http://biotech.about.com/od/pcr/a/PCRtheory.htm

http://www.carolina.com/pcr-kits/dnalc-human-mitochondrial-dna-extraction-and-amplification-kit-with-05-ml-tubes/211236.pr

http://chemistry.about.com/od/chemistryterminology/a/What-Is-A-Control-Group.htm

http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA460885

http://www.ehow.com/way_5255896_pcr-basics.html

http://faculty.ccbcmd.edu/courses/bio141/labmanua/lab17/lab17.html

http://www.genome.gov/10000207

http://medical-dictionary.thefreedictionary.com

http://www.nfstc.org/pdi/Subject02/pdi_s02_m02_01_b.htm

http://www.nfstc.org/pdi/Subject01/pdi_s01_m02_01.htm

http://www.niaid.nih.gov/topics/immunesystem/Pages/whatIsImmuneSystem.aspx

http://projects.nfstc.org/fse/01/01-05.html