Cancer Cell Technology and Immunology Principles

Robert Aguilar

ABSTRACT:

Testicular cancer mainly affects men between the ages of 20 and 39, but it is the most common neoplasm in men between the ages of 15 and 34. The National Cancer Institute states that localized testicular cancer has a recurrence rate of 15-20%, and those tumors that are determined to be Sertoli or Leydig cell derived do not respond well to chemotherapy or radiation treatment. My summer research was centered on the effort to better characterize the immune response to a testicular tumor. This was accomplished through the use of different immunology techniques such as proliferation assays, immunohistochemistry, ELISA, and ELISPOT. My lab experience inspired me to design a unit curriculum which will help elucidate the various physiognomies of tumor growth. Students first learn and apply the essential techniques of cancer cell culture. The testicular cancer cell line I-10 (CCL-83, ATCC) is used to familiarize students with routine feeding and maintenance, subculturing, counting, and cryopreservation. I-10 cell RNA is converted to cDNA and qualitatively analyzed for inhibin-7ap0(g)10(a)4(nd22s2(ne)c2(ne)c)Tj tiompee wiTJ si,3(i)-2-2(t)-2(ur)20(z)-RTTd [(t)1c 0.004 TTJ 3.5

Autoreactive T Cells

T cells recognize antigens only when presented by molecules on the surface of antigen presenting cells such as macrophages and dendritic cells[14]. CD4+ T cells recognize peptide antigens via the T cell receptor (TCR) when the peptide is presented by class II molecules of the major histocompatibility complex (MHC). However, CD8+ T cells recognize peptide antigens when they are presented by class I molecules of the MHC. In most organ-specific autoimmune models, CD4+ responses predominate, but CD8+ T cells may be involved in the self-recognition process and in the pathogenesis of disease often by recognizing totally different peptide domains of self-proteins than those recognized by CD4+ T cells[15].

STUDENT OUTCOMES:

Students learn and apply the essential techniques of cancer cell culture. The testicular cancer cell line I-10 (CCL-83, ATCC) is used to familiarize students with routine feeding and maintenance, subculturing, counting, and cryopreservation. I-10 cell RNA is converted to cDNA and qualitatively analyzed for inhibin-alpha gene expression, which serves as a tumor marker, using RT-PCR. Then, they analyze previously formalin-fixed, paraffin embedded murine testes (normal and cancerous) using H&E staining and immunohistochemical staining for CD3 and inhibin-alpha. Finally, students complete this unit bynd(us)3(e)-3(r)-5(7 0 Td 0 nd1 m-a(k)-2(s)(ul)-1(t)-3(3)-2(7)-1(u)-2(0)-2(n)-2(d)-2(s)Tc 0.00

ADVANCE PREPARATION:

Materials	Purchased	Cost	Substitute
I-10 Leydig Cancer Cells (CCL-83)	ATCC	$450	N/A
F-12K Medium (30-2004)	ATCC	$18	CO_2 Independent Medium
Horse Serum (30-2040)	ATCC	$50	N/A
Fetal Bovine Serum (SH3091002)	Fisher Scientific	$70	N/A
Penicillin/Streptomycin (BP2959-50)	Fisher Scientific	$50	N/A
Trypan Blue (SV3008401)	Fisher Scientific	$20	N/A
Vacuum Filtering Flask (10-182-50B)	Fisher Scientific	$30	N/A
Cell-VU Hemocytometer Slides (DRM700)	Fisher Scientific	$200	N/A
Permount			

Sterile Materials

1. The following items should be sterilized by spraying with 70% ethanol and left permanently inside the cell culture hood (laminar flow hood):
    a. Permanent marker
    b. Test tube racks (Do not use wood!)
2. The following items should be sterilized by steaming and left permanently inside the cell culture hood (laminar flow hood):
    a. Pasteur pipettes (in pipet sterilizing box)

Horse Serum (HS)

1. *Time required: Thaw-1.0 hr., aliquot-30 min.*
2. Thaw HS by placing it in a 37ºC water bath.
3. Aliquot into 50mL conical tubes.
4. DO NOT decomplement!
5. Store at -20 C.

Fetal Bovineliqu2 1 Tf0 h.5,0 Tliqud(T)1(e)4(s)-1(t)-2(ms)-1(t)-2(ms)-1(t)-2(

MATERIALS AND EQUIPMENT

Student Workstation Checklist

One workstation serves 2-3 students.

Cell Subculture (Cell Culture Hood)		
Item	**Contents/Volume**	**Number ( )**
Inverted Microscope		1
CO_2 Incubator		1
Water Bath		1
Clinical Centrifuge		1
Gloves		2-3
Beaker (for ice)		1
Ice	600mL	1

Cryopreservation of Cancer Cells
(Cell Culture Hood)

Handling Procedures for Frozen Cells (Cell Culture Hood)		
Item	**Contents/Volume**	**Number ( )**
Inverted Microscope		1

Cell Counting **Trypan Blue Exclusion (at bench)**		
Item	**Contents/Volume**	**Number ( )**
Inverted Microscope		1

DAILY UNIT PLANS

Thawing Procedure for Direct Cell Shipment from ATCC (Teacher)

1. This should be started at least **1 week** in advance in order to ensure that you have enough cells for all of your student workstations.
2.

19. Once you have a single cell suspension, pipette 10mL of the **RINSE** tube into one culture flask.
20. Now using a new serological pipette, add 3mL of new F-12K complete medium to each flask in order to bring the volume up to a total of 13mL.

Subculturing Cells (Week 1, Days 1, 2, 4/Week 2, Days 1 & 4/Week 3, Day 1)

1. At least 30 minutes before you begin, turn on the 37°C water bath and place F-12K complete growth medium inside to warm it and the TE for thawing.
2. **NOTE:** You will need 5mL of TE for each culture flask.
3. Sterilize your gloved hands by squirting 70% EtOH on them. Then rub your gloved hands together as if you are washing them.
4. Obtain your cell culture flasks from the CO_2 incubator, and take a look at them under the inverted microscope. Healthy growing I-10 cells will have adhered to the bottom of the flask, and some should have a spindle shape. See below.

5. Estimate percent confluency by comparing the amount of space covered by the cells with the unoccupied spaces. The figure above is about 50% confluent. Once you have about 80% confluency and most cells are spindle-shaped, they need to be subcultured.
6. **NOTE:** In order to prevent contamination, never place the culture flask directly on the bench top. Instead, first place a clean paper towel underneath the flask.
7. Obtain a sterile 50 mL tube, and use the marker that is already in the cell culture hood to label it "**SUB**, **Station #and/or initials**" (one 50mL conical tube for each flask).
8. Now, unscrew the cap on the culture flask while inside the cell culture hood, and place the cap on the floor of the cell culture hood so that the open end faces up.
9. Using the supernatant from within the culture flask, rinse the "belly" of the flask by squirting it with the supernatant in a side to side motion. Do this about 5 times. This will help initiate the dislodging process.
10. **NOTE:** It is important not to let the cells sit dry (without growth medium) for too long otherwise their viability will be compromised.
11. Pipette loosened cells from flask into the corresponding **SUB** tube.
12. Place tube on ice.
13. **DO NOT** let the cells in the flask sit dry for too long!
14. Squirt the TE tube with 70% EtOH to sterilize it before placing it inside of the cell culture hood. Pipette 5 mL of TE into the 25cm2 flask (10 mL if using a 75cm^2 flask).
15. Carefully, place flask in CO_2 incubator for at least 3 minutes in order to dislodge cells from the flask. (Repeat for other flasks.)
16. Remove your TE/cells/flask from the CO_2 incubator and again pipette TE/cell solution from within the flask, and use it to rinse the "belly" of the flask by squirting it from side to side. Do this about 5 times.

32. Dispose of any used tubes, pipettes or flasks that have come into contact with the cells in the biohazard bin.
33. Wipe the floor of the cell culture hood with 70% EtOH to sterilize.
34. Make sure that there are no perishables such as cells or culture media still inside the cell culture hood.
35. Close the hatch and turn on the UV light.

Cryopreservation of Cancer Cells (Week 3, Day 1)

1. This procedure is for culture flasks that are at about 80% confluency. I generally do not let students go beyond 4 flasks.
2. At least 30 minutes before you begin, turn on the 37°C water bath and place F-12K complete growth medium inside to warm it and the TE for thawing.
3. **NOTE:** You will need 5mL of TE for each culture flask.
4. Sterilize your gloved hands by squirting 70% EtOH on them. Then, rub your gloved hands together as if you are washing them.
5. Obtain your cell culture flasks from the CO_2 incubator, and take a look at them under the inverted microscope. Healthy growing I-10 cells will have adhered to the bottom of the flask and some should have a spindle shape. See below.

6. Estimate percent confluency by comparing the amount of space covered by the cells with the unoccupied spaces. The figure above is about 50% confluent. Once you have about 80% confluency and most cells are spindle-shaped, they can be cryopreserved.
7. **NOTE:** In order to prevent contamination, never place the culture flask directly on the bench top. Instead, first place a clean paper towel underneath the flask.
8. Obtain a sterile 50 mL tube and use the marker that is already in the cell culture hood to label it "**FROZ**, **Station #and/or initials**" (one 50mL conical tube for each flask).
9. Now, unscrew the cap on the culture flask while inside the cell culture hood, and place the cap on the floor of the cell culture hood so that the open end faces up.
10. Using the supernatant from within the culture flask, rinse the "belly" of the flask by squirting it with the supernatant in a side to side motion. Do this about 5 times. This will help initiate the dislodging process.
11. **NOTE:** It is important not to let the cells sit dry (without growth medium) for too long otherwise their viability will be compromised.

Cell Counting, Trypan Blue Exclusion (Week 2, Days 1 & 4)

1. At least 30 minutes before you begin, turn on the 37°C water bath and place F-12K complete growth medium inside to warm it and the TE for thawing.
2. **NOTE:** You will need 5mL of TE for each culture flask.
3. Sterilize your gloved hands by squirting 70% EtOH on them. Then, rub your gloved hands together as if you are washing them.
4. Obtain your cell culture flasks from the CO_2 incubator and take a look at them under the inverted microscope. Healthy growing I-10 cells will have adhered to the bottom of the flask and some should have a spindle shape. See below.

5. Estimate percent confluency by comparing the amount of space covered by the cells with the unoccupied spaces. The figure above is about 50% confluent. Once you have about 80% confluency and most cells are spindle-shaped, they need to be subcultuted.
6. **NOTE:** In order to prevent contamination, never place the culture flask directly on the bench top. Instead, first place a clean paper towel underneath the flask.
7. Obtain a sterile 50mL tube and use the marker to label it with your initials and the cell culture you will be working with.

17. Pipette the TE/cell solution from the flask into the corresponding **SUB** tube with cells that you had on ice. (Repeat for other flasks.)
18. Place **SUB** tubes in centrifuge and spin @ 1,500 rpm, 4°C, for 5 minutes.
19. While the tubes are spinning, get as many new culture flasks as you need and label them with "**Cell type, Station #,** and **Date** inside the cell culture hood. For every **SUB** tube, you will need 2 flasks.
20. After the spin, the **SUB** tubes at this point can either be left in the centrifuge @ 4°C while you work or can be put back on ice. Be careful not to disturb the cell pellet while you are handling it!!! If you do, you must spin it down again for 5 minutes.
21. Squirt the outside of the **SUB** tube with 70% EtOH to sterilize it, and then, place it in the cell culture hood. , and then,

33. Keep track of your cell count using a Lab Counter App.
34. Pipette 5mL of the **SUB** tube into one culture flask and 5mL into the other culture flask.
35. Now, using a new serological pipette, add 2mL of F-12K complete medium to each flask in order to bring the volume up to a total of 7mL.
36. Take a look at the cell-filled culture flasks under the inverted microscope.
37. At this point, all of the cells will be suspended in the medium and should look relatively round and dense. See below.

38. Now, say a short prayer and carefully place it in the CO_2 incubator. Culture flasks may be stacked.
39. Dispose of any used tubes, pipettes or flasks that have come into contact with the cells in the biohazard bin.
40. Wipe the floor of the cell culture hood with 70% EtOH to sterilize.
41. Make sure that

10. Carefully, pull comb straight up without wiggling.
11. If you notice "dimples" from the gel in the buffer, slowly add more buffer until they are completely submerged.

Load Gel

12. Add 1µL of loading dye to your PCR reaction and mix by pipetting up and down.
13. Load a 1kbp DNA Ladder into the first well and then 20µL of your PCR reaction into each well after.
14. Draw a gel map that shows which wells were loaded with what sample.
15. Connect to the power supply and set it for 150V for 30 minutes.
16. Unplug chamber and view gel on Transilluminator.

Hematoxylin & Eosin Staining (Week 1, Day 3)

1. Tissue samples were obtained, embedded in paraffin wax, and sliced at the Cleveland Clinic.
2. Place slides containing paraffin sections in a slide holder (glass or metal).
3. The following steps should be completed under a fume hood.
4. Deparaffinize and rehydrate sections as follows:
    a. 3 x 3 min. 100% Xylene
    b. 3 x 3 min. 100% Ethanol
    c. 1 x 3 min. 95% Ethanol
    d. 1 x 3 min. 80% Ethanol
    e. 1 x 5 min. dH_2O
5. Hematoxylin staining:
    a. 1 x 3 min. Hematoxylin
    b. Rinse dH_2O
    c. 1 x 5 min. tap H_2O
    d. Dip 8-12x (fast) Acid Ethanol (to destain)
    e. Rinse 2 x 1 min. tap H_2O
    f. Rinse 1 x 2 min. dH_2O (can leave overnight at this stage)
6. Blot excess water from the slide holder before going into Eosin.
7. Eosin staining and dehydration:
    a. 1 x 30s. Eosin
    b. 3 x 5 min. 95% Ethanol
    c. 3 x 5 min. 100% Ethanol
    d. 3 x 15 min. Xylene
8. You can leave slides in xylene overnight to get good clearing of any water.
9. Coverslip slides using Permount (xylene based).
    a. Place a drop of Permount on the slide using a micropipette, taking care not to leave any bubbles directly over the tissue.
    b. Angle the coverslip, and let it fall gently onto the slide.
    c. Allow the Permount to spread beneath the coverslip, covering all the tissue.
    d. Dry overnight in the hood.
    e. In case any of the Permount spills over the slide, it's a good idea to set the finished slide on a plastic platform for drying and not a paper towel (pipette tips box works well). Otherwise, the paper towel will be permanently glued to the slide.

Immunohistochemistry (Week 2, Days 2 & 3)

Rehydrate Paraffin Embedded Tissue (30 min.)

All steps should be completed in a Coplin Jar on gentle shaker within a fume hood.
Some of these steps will be completed using the ImmunoCruz Staining System.

1) Xylene 2 x 5 min @ R.T.
2) 100% EtOH 1 TfqT80CID 7 >> R.T.

2)95j /TT1 1 Tf 0.67 0 Td [(1y)20(l)-2(e)4(n)-10(e)4(2)0(l TfqT80CID 7 >> R)-3(.T)1(.)]TJ 10.14

24) Counterstain slides in Gill's formulation #2 hematoxylin for 5-10 seconds. Immediately, wash with several changes of deionized H_2O.
25) Dehydrate sections as follows: 2x 70% ethanol for 10 seconds each, 2x 95% ethanol for 10 seconds each, 2x 100% ethanol for 10 seconds each and 3x xylenes for 10 seconds each. Wipe off excess xylene from edges and bottom of slide.
26) Immediately, add 1–2 drops of permanent mounting medium directly on the tissue and cover with glass coverslip.
27) **DO NOT LET XYLENE DRY OR THE TISSUE WILL CRACK.**

AP Biology Lab Practical

Lab Bases

Direction

Molecules

2)