Sounding the Alarm: Studying the Role of Signal Cascades in the Immune Response

Mr. Mark Trice
Frederick High School
650 Carroll Parkway
Frederick, MD 21701

Mark.trice@fcps.org

Funded by: The American Association of Immunologists
Mentored by David Rozak, Ph. D
United States Army Medical Research Institute of Infectious Diseases (USAMRIID)
Frederick, MD 21701

Abstract

Sounding the Alarm: Studying the Role of Signal Cascades in the Immune Response.
Mark Trice[1] and David Rozak, Ph. D.[2]
[1] Frederick High School, Frederick, MD [2] United States Army Medical Research Institute of Infectious Diseases (USAMRIID), Frederick, MD

The purpose of this research opportunity was to investigate the methods and roles of cell signaling via protein cascades and their connections to the functionality of the immune system. During my summer internship, I worked with investigators at the United States Army Medical Research Institute for Infectious Diseases (USAMRIID) to develop and test the activity of man-made protease cascades that could someday mirror the complement system in its aggressive response to specific signal molecules.

My research experience aimed at understanding the role and function of protein cascades in immune responses inspired me to develop a unit for a second year Biology/AP Biology course. Students examine the effects of several factors, including protease concentration, substrate concentration, and a non-cascade reaction, on reaction rates through construction of both physical models and simulated protease cascades. In this unit, students first work with a hands-on physical, manipulative model to familiarize them with how a protein cascade works. Then, all of the students will work in tandem on a kinesthetic activity modeling exactly how protein cascades behave.

This activity allows for a cost-effective method of demonstrating protease cascades without expensive lab equipment and reagents. Students translate their knowledge into the application stage when they examine how protease cascades work in the human body and discuss various uses and effects of cascades, such as protein kinases, on the immune system.

Instructor Section

Science Background

These mechanistic cascades can be used to create a programmable molecular sensor which would respond to an analyte[11] and subsequently amplify and measure the signal of the reaction. The onset of the signal can then be correlated with the concentration of the initiating analyte.

What will students do?
Students will utilize a physical, manipulative model to demonstrate the pathway of a protease cascade. With background knowledge and this model, students will take part in creating a kinesthetic representation of a protease cascade, namely the subtilisin cascade. During the simulation, students will collect data and graph their results to conclude the effects of several variables on reaction rate. Students will then apply their knowledge to additional comprehension and application assessments that allow them to explain the model in their own words.

Placement
This unit is placed in the cellular signaling and pathways chapter. Students are already familiar with biochemistry applications, such as enzyme kinetics, and are also familiar with the types of cells and cell communication. The concepts are also reiterated in the chapter about immunology within the human body unit in the AP Biology class.

Student Outcomes
- Students will be able to model and explain what a protein cascade is and how it works.
- Students will be able to explain the interplay between enzymes and active sites, and how these interactions affect protein cascades.
- Students will model the release of the prodomain from the subtilisin complex and physically model how the cascade works.
- Students will explain the importance of fluorescence in the protein cascade and how these molecules allow for measurement of reaction rate.
- Students will construct, analysis, and extrapolate information from graphs to determine conclusions when various parameters are manipulated.

[10] Lunder M, Bratkovic T, Kreft S, Strukelj B (July 2005). "Peptide inhibitor of pancreatic lipase selected by phage display using different elution strategies". J. Lipid Res. **46** (7): 1512–6
[11] **Analyte- substance undergoing analysis**. "Analyte." Merriam-Webster. Merriam-Webster, n.d. Web. 10 July 2013. <http://www.merriam-webster.com/medical/analyte>.

Time Requirements

This unit is designed for a 90 minute block period for a 6 day period with an additional day for project presentation. Below is the unit schedule:

Day Number

Lesson Plans

Day 1: Introduction to Cell Signaling and Communication

Engagement:

Students will view the following two videos, one of dominoes falling, and the other on Rube Goldberg machines. After watching the video, brainstorm on the board what the similarities are between the two videos. Write all answers on the board, and then direct the conversation to the fact that both have a "cascading" effect, where one object sets off a chain reaction.

- Domino/Rube Goldberg analogy
  https://www.youtube.com/watch?v=dLRjiiAawGg
  https://www.youtube.com/watch?v=uF3nV0r87v8

Elaboration:

Students will view the following video on general cell signaling. Have them take notes and/or write down any important information from the video.

YouTube Video: Bozeman Biology: Cell Signaling:
https://www.youtube.com/watch?v=qOVkedxDqQo

Explanation:

Students will read the cell signaling article on cellular signaling. They will underline all important information as they read.

Exploration:

Students will then complete the reading comprehension guide about the cell signaling article.

Evaluation:

As a class, review the main ideas of the article and how to use the reading guide to facilitate comprehension.

Day 2: Protease Cascade manipulative and "Tag- You're Lit"

Explanation:

Students will each receive a bag with the manipulative pieces already cut out and ready. Follow directions on the paper and as a class, simulate a protease cascade.

Engagement:

Students will receive a paper copy of the "Tag-You're Lit" lab paper. Students will be given roles and act out each of the three trials of the activity.

Evaluation:

Students will collect their data and graph the data to show trends.

Exploration:

Students must discuss in their group what trends exist, what the trends in the data suggest, and how that relates to real-life protease cascades

Day 5: Erectile Dysfunction Group Activity

Engagement:

As a class, all “Fire!” analogies are presented, and students are asked to vote for their favorite construct that helps to parallel the protease cascade.

Day 6: Group Activity and Immunology presentation introduction**

Explanation:

Students will continue to work on the group activity and share their answers with other groups.

Engagement:

Students will receive a copy of the immunology/cell signaling activity and begin to brainstorm ideas. By the end of class, students must present to the teacher what disease they would like to work with and a rough outline of their important points/main ideas they want to cover. They are allowed to use computers for the entire class.

**this activity can be continued for however long you'd like to give for research

Teacher Version: Cell Signaling

What Kind of Signals Do Cells Receive?

Cells must be able to receive signals, as well as propagate signals to elicit a response. Chemical in nature, cell signals can be used as sensor mechanisms, growth factors, hormones, and neurotransmitters. Neurotransmitters, a class of short-range signaling molecules that travel across the neuromuscular junction, aid in intracellular signaling among neurons. Other molecules, such as FSH (Follicle Stimulating Hormone), move further from the targeted cell traveling from the brain to the ovary, resulting in egg release form the ovary. Other signaling molecules aid in mechanical stimulation of cells, such as cells in the ear reacting to movement of sound waves, baroreceptors that detect changes in blood pressure in the vascular system, or pressure sensors in the skin.

How Do Cells Recognize Signals?

Extracellular receptor proteins bind to signaling molecules in order to initiate certain types of responses, whether they are chemical or physiological. There exist different types of receptors, depending upon their target and/or function; dopamine receptors bind to dopamine, insulin receptors bind to insulin, etc. The signal pathway begins when transmembrane protein receptors bind to extracellular signaling molecules and subsequently transmit the signal through a series of "molecular switches" which in turn initiates signaling pathways within the cell resulting in a cell signal being propagated within the cell. The membrane receptors that initiate the intracellular signals include three different classes: G-protein coupled receptors, ion channel receptors, and enzyme-linked receptors.

For example, gated ion channels, such as acetylcholine receptors allow for soluble materials to travel across the membrane when the pore is open, and, when the pore is closed, no external signal is present. Therefore, when an acetylcholine molecules bind to a receptor, a conformational change within the membrane allows for transport of ion is through the membrane.

How Do Cells Respond to Signals?

After the signal is received from the receptor protein, a conformational change occurs, thus launching series of biochemical reactions known as a signal transduction cascade. This amplifies the message and propagates the signal throughout the cell. After the receptor activates the cascade, secondary messengers are triggered which initiate and coordinate the intracellular signaling pathways. The quintessential example is cAMP, a common secondary messenger which contributes the increasing propagation of the signal. cAMP, synthesized from

Teacher Version: Tag- You're Lit!

The goal of this activity is to show how reaction rate and fluorescence varies when different parameters of the protein cascade reaction are manipulated.

ATTENTION!

For all simulations, you will need to:

1. **Choose one person as a "student timer" that will sit at the "fluorescent end zone" and record the time from when the simulation started until each person crosses the line into the designated area.**
    a. **Alternatively, the teacher could set up a video camera that would run for the entirety of the simulation. This provides the opportunity for all students to participate as well as provides a digital record of data.**
2. **Choose one student as the "free subtilisin" (FS) molecule that will begin the simulation by tagging one of his/her fellow students, thus beginning the reaction.**

Simulation 1: Reaction Rate

Here the speed with which a person gets up from a sitting position will show how reaction rates can vary

Trial 1:

1. Spread out all students on a large field, i.e. football or soccer field, or large gymnasium. These students should be sitting on the ground.
2. Hand out a "secret card" to all persons playing, EXCEPT the FS student. These cgSord of data.

3. Cell signaling

6. Describe what would occur if a cell has activated protein kinase enzymes that are activated over a long period of time.
    a. **<u>The transcription factor will continually transcribe the DNA strand, given that the protein kinases remain active.</u>**
7. What would occur in a cell if the protein kinase enzymes were deactivated?
    a. **Phosphate groups**

Name(s) _____KEY_____

Block _____ Date _____

Cell Signaling/Immune System Group Quiz

1. Describe the three lines of defenses we as humans have against germs. Also explain how each line protects us.

The three lines of defense include the first, second, and third line of defense. The first line of defense is the external barriers, such as skin or mucous, that serves as the protection nonspecifically against all pathogens and invaders. The second line of defense is characterized by an inflammatory response, where, upon introduction of a foreign object, such as a splinter or needle, the body mounts a defense. Blood circulation is increased to the infected site via vasodilation after histamines are released. Next the body releases phagocytes, such as neutrophils or monocytes, to block further invasion by viruses. Lastly the body mounts the third line of defense, which is very specific. The body produces B lymphocytes which produce antibodies to the infecting particle. The body will use this to fight the infection, however the body will also make memory cells to "remember" this invader and mount an attack if it ever encountered it again. T lymphocytes are also released in a cell-mediated response. This is a type of "first-hand combat" where the T cells activate macrophages which go and engulf the particle and destroy it. This type of response also relies on antigens on cells to elicit a specific response to that specific molecule.

2. Define a signal transduction pathway
    a. **In signal transduction pathways, a single molecule can stimulate the release of thousands of molecules of product within a cell.**

3. Consider several examples of intercellular signaling, such as kinase pathways. How does a protein kinase cascade amplify an intercellular signal?
    a. **One activated receptor kinase molecule can trigger the activation of thousands of proteins.**

4. Nur77, an orphan nuclear receptor, plays a key role in apoptosis in T cells. It was also shown in an experiment by Wang et al that Nur77 is the physiological substrate of the MEK-4(gn a)4(91648.90 Td (4. 514)-1e(re)-1t)-2(r)3(a)

Phosphorylation cascades involving a series of protein kinases are useful for cellular

Student Section

Adapted from:

Cornell Notes Template



Tag- You're Lit!

The goal of this activity is to show how reaction rate and fluorescence varies when different parameters of the protein cascade reaction are manipulated.

ATTENTION!

For all simulations, you will need to:

1. **Choose one person as a "student timer" that will sit at the "fluorescent end zone" and record the time from when the simulation started that each person crosses the line into the designated area.**
    a. **Alternatively, the teacher could set up a video camera that would run for the entirety of the simulation. This provides the opportunity for all students to participate as well as provides a digital record of data.**
2.

Trial 2:

1. Gather your student timer as noted before.
2. This time, three students will become FS students. Each of them will proceed to the endzone as in Trial 1, tagging others along the way.
3. Record times that each person entered the endzone and broke their lightstick to show fluorescence.

Trial 3:

1. Repeat the above procedure, however this time, start with five FS students.
2. Gather data and record times that each person entered the endzone and broke their lightstick to show fluorescence.

Simulation 3: Non-cascade Reaction

In this simulation, some students will become uninhibited and be able to proceed with the cascade, while others, once taggedbc(s)4(t)2(2 72 44(t)2(2 724nc)14(e.)t)2 /P <<24ncy10(wgoc

Name (s) _____

Erectile Dysfunction and the Cell Signaling Pathway

Cell signaling can have real-life applications, one common problem among men being erectile dysfunction. While many medications can be prescribed, a holistic approach has been suggested among the medical community.

Within your group, complete the following tasks and be prepared to explain your research to other groups in the class.

1. Research what erectile dysfunction is, what causes it, and how it is currently treated.
2. Examine the cell signaling pathways that exist in this disorder. Be sure to address topics such as the primary messengers of nitric oxide and cGMP as a secondary messenger, as well as how all these events correspond to interactions with Ca++ and myosin to produce an erection.
3. Address the holistic approach of administering ginseng and how this mechanism works. What effects does ginseng have on nitric oxide levels? What effects does Viagra have on PDE5 and cGMP levels in the body?
4. Provide your recommendations as the patient's doctor as to what course of treatment you find most effective and what specific dosages would you prescribe?

Name(s) _____

Phosphorylation Cascade

1. Draw a cell signaling cascade. Now show specifically when phosphorylation occurs and

IMMUNOLOGY AND CELL SIGNAL PROJECT

Grading: Total Points - 80

Format - poster - 15 points

Traditional Poster

- x
- x

Name(s) _____
Block _____ Date _____

Cell Signaling/Immune System Group Quiz

1. Describe the three lines of defenses we as humans have against germs. Also explain how each line protects us.

2. Define a signal transduction pathway du(f7mC /LTd( )TjEMC /P <MCID 10 >27(w)-5(a)-

   2.

4. Nur77, an orphan nuclear receptor, plays a key role in apoptosis in T cells. It was also shown in an experiment by Wang et al that Nur77 is the physiological substrate of the MEK-ERK-RSK cascade, which functions as a phosphorylation cascade for the intracellular translocation during T cell death.

   Wang, A, J Rud, CM Olson, J Anguita, and BA Osborne. "Phosphorylation of Nur77 by the MEK-ERK-RSK cascade induces mitochondrial translocation and apoptosis in T cells.." Immunology 183: 3268-77. Print.

   Phosphorylation cascades involving a series of protein kinases are useful for cellular signal transduction because
   a. They are species specific
   b. They always lead to the same cellular response
   c. They amplify the original signal manyfold
   d. They counter the harmful effects of phosphatases

5.

Matching: Match the word with the description below. You may use words more than once, once, or not at all.

B cells	Macrophages
Helper T cells	Cytotoxic T cell

7. Bonds to class II MHC molecules _____
8. An APC that is also a part of innate immunity_____
9. Responsible for humoral immunity _____
10. A lymphocyte that secretes cytokines to stimulate other lymphocytes _____

11. Describe three responses by the innate immune system in response to a pathogen.

12. Explain the positive feedback loop in regards to how Helper T cells use cell signaling to initiate an active immune response.