AAI High School Teachers Summer Research Program in Immunology:

I. Science Background **3**
Primer of Immunology **3**
Brief Review of the Immune System 3
Dengue Q & A **5**
Glossary of Basic Immunology Terms **6**
II. Student Outcomes 7
Concepts Covered: **7**
Standards Addressed: **7**
Placement in Course **8**
What students will do and technical skills learned **8**
III. Learning Objectives 8
IV. Time Requirements 8
V. Advance Preparation 8
VI. Materials and Equipment 8
VII. Student Prior Knowledge. 8
VIII. Daily Unit Plans 9
Daily Lesson Plans **9**
IX. How will students demonstrate understanding? 11
X. References 11
XI. STUDENT SECTION 12
..... **13**
What are PCR, RT-PCR and Flow Cytometry? **16**
Dengue - Is the World Cup Worth the Risk? **19**
XII. APPENDIX 25

when a T or B cell is activated through antigen binding causing it to divide to form clones (genetically identical cells). This is called clonal selection.

iii. Distinguishing Self from Nonself – Through clonal deletion B or T cells that have the potential of attacking one's own body cells are eliminated through apoptosis.

There are five classes of antibodies secreted by B cells.

IgM – the first class of antibodies produced after the initial exposure to an antigen and is short lived.

IgG – is the most abundant antibody in blood and tissue fluids.

IgD – acts as cell surface receptor on B cells.

IgA – provides protection for mucus membranes.

IgE – present in blood, triggers release of histamines and other chemicals involved in allergic reactions.

Dengue Q & A

Question: What is dengue?
Answer: Dengue is a vector borne viral disease caused by one of four different serotypes (DENV-1, DENV-2, DENV-3, DENV-4). The virus is a member of the Flaviviridae family.

Question: How is it contracted/transmitted?
Answer: The most common vector is the *Ades aegypti* mosquito. *Ades albopictus* is also a carrier. When an infected mosquito bites a human, the virus can be transmitted through the fluids the mosquito inserts through its proboscis. The mosquito acquires the virus through the blood of an infected human it bites. Human to human transmission is not possible.

Question: Can I get dengue again if I have already been infected?
Answer: You will have immunity to the specific virus (DENV-1, DENV-2, DENV-3, DENV-4) that you were originally infected with and, if infected again with that serotype, you will not get the disease. However, if you are infected with a different serotype you are at a greater risk of acquiring dengue hemorrhagic fever (DHF). DHF is a more severe form of dengue and can be fatal if medical treatment is not immediately sought.

Question: How can it be prevented?
Answer: Because the disease is spread by mosquitoes, eradication of breeding sites is essential. *Ae. aegypti* lays her eggs in containers that hold water (e.g. tires, planters, water drums). However, *Ae. aegypti* are adaptable to varying conditions; eggs have been known to survive drought conditions. *Ae. albopictus* prefers cool dark areas such as home closets. The use of air conditioning and screens can prevent adults from finding their human meals.

Question: What are the disease symptoms?

Question: How is dengue treated?
Answer: There is no specific treatment for the disease. If one suspects they have dengue fever, they can take analgesics, stay hydrated, rest, and speak with their doctor. If they develop more severe symptoms after the fever has subsided, they should go to a hospital. Fluid replacement therapy is appropriate at this point. Currently, there is no vaccine on the market; however the company Sanofi Pasteur has plans to register the tetravalent vaccine (protects against all 4 dengue strains) in 2015 with a target distribution time frame of late 2015.

Questions: Where has the disease been found?
Answer: The four viruses originated in monkeys and spread to humans in Africa or Southeast Asia between 100 and 800 years ago. Currently, the disease can be found in Southeast Asia, the Pacific Islands, Africa, Caribbean, and the Americas.

Question: What lab tests are used to detect the virus?
Answer: The test methods used is dependent upon the duration of the disease.

Real time RT-PCR is used to detect the virus in patients within the first 5 days of symptoms. A positive result confirms presence of the disease and usually which viral strain is involved. A negative result in this time frame is termed indeterminate, and patients will be asked for another sample 5 days post symptoms.

MAC ELISA detects the presence of IgM antibody in the serum. The test can be used as early as 3 days post symptoms and may be able to detect the antibody as far as 90 days post symptoms. Caution should be used interpreting the results as this method can also react with other flaviviruses.

IgG ELISA is used to diagnose the presence of a past infection. To determine if the infection

e)13(d to)ph)14(o)2(s)-7(i)-7(t)-8(i)-8(v)-11(e)-11(.)-11(T)-11(h)-8(e)-11(A)-11(n)-8(t)-11(i)-8(g)-11(e)-11(n)-8(-)-11(s)-8(p)-11(e)-11(c)-8(i)-11(f)-11(i)-8(c)-11(.)-11(T)-11(h)-8(e)-11(I)-11(g)-8(M)-11(.)-11(T)-11(h)-8(e)-11(I)-11(g)-8(G)-11(.)-11(T)-11(h)-8(e)-11(I)-11(g)-8(A)-11(.)-11(T)-11(h)-8(e)-11(I)-11(g)-8(D)-11(.)-11(T)-11(h)-8(e)-11(I)-11(g)-8(E

Complement Proteins - a set of diverse proteins that function in a cascade manner. One complement protein binds to proteins on the pathogen which alerts phagocytes to the presence of an invader. Another protein activates the inflammatory response. Other proteins are responsible for lysing the pathogen.

Cytokines – signaling proteins that affect the behavior of their target cell.

Immunity – the body's ability to resist disease when invaded by a pathogen.

Inflammation – damaged tissue that exhibits puffiness, warmth, and redness.

Interferon – a class of cytokines produced by virally infected cells. This signal protein alerts neighboring cells increasing their ability to resist infection.

Lymphocyte – type of white blood cell that is in involved in adaptive immunity; it includes B cells and T cells.

Major Histocompatibility complexes (MHC) – proteins found on a host cell's surface that hold an antigen fragment that can bind to a T cell receptor.

Natural Killer Cells – a type of lymphocyte that recognizes cells that are infected or cancerous and initiates cell destruction. They are part of the innate immune system.

Pathogen - organisms and viruses that can cause disease when they infect a host.

Phagocyte – white blood cells that engulf pathogens, they are involved in innate and adaptive immunity (e.g. macrophage).

T cells (T lymphocytes) –mature in the thymus. They have a variety of functions: activating macrophages, helping B cells produce antibodies, and killing infected cells.

T cell receptors – proteins found bound in the membrane of T cells. They recognize and bind to non-self substances (antigens) bound to MHC proteins on other cells.

II. Student Outcomes

Concepts Covered:

Students will have a deeper understanding of the adaptive immune system. Through the use of virtual labs and videos, students will be familiarized with the techniques of ELISA, PCR, RT-PCR, and flow cytometry. By completing the dengue case study, students will learn about the cause, symptoms, diagnosis, and treatment of dengue fever and dengue hemorrhagic fever.

Standards Addressed:

AP Biology

Essential knowledge 2.D.4 – Plants and animals have a variety of chemical defenses against infections that affect dynamic homeostasis.

Essential knowledge 3.D.2 – Cells communicate with each other through direct contact with other cells or from a distance via chemical signaling.

Next Generation Science Standards

HS-LS1-2. Develop and use a model to illustrate the hierarchical organization of interacting systems that provide specific functions within multicellular organisms.

HS-LS1-3. Plan and conduct an investigation to provide evidence that feedback mechanisms maintain homeostasis.

Placement in Course
These lessons can be incorporated in an Immunology unit or Biotechnology unit. The lessons are designed for an Honors Biology and/or AP Biology course.

What students will do and technical skills learned
Students will review/learn the basics of the adaptive immune system. Through the use of cell images, students will reinforce their knowledge of the various forms of B and T cells and how they interrelate. Students will complete a virtual ELISA

VIII. Daily Unit Plans

Daily Lesson Plans

General overview of unit - The unit is designed to introduce or reinforce the concepts of the adaptive immune system to students using dengue fever as a real-life application.

Assign the video and reflection guide (see Student Section) to be completed prior to the start of the unit.
http://www.bozemanscience.com/immune-system

Day 1 – Teacher will review or introduce (depending on the class level) the components of the adaptive immune system; how different cell types interrelate and the general schemata of how the human body responds to a pathogenic insult.

> *Activity* -could be used Day 1 as reinforcement or Day 2 as formative assessment. Each student will be given a set of images that contain the different cell types of the adaptive immune system. The images are located in the Student Section. Students will first be asked to separate them into different cell types (e.g. B and T cells). Then, using the cell images students will create a flow chart representing how the immune system responds to an exposure to a pathogen

c) Reverse Transcription PCR

Prevention

This Blog discusses the use of genetically modified mosquitoes to control their population size.
http://www.nature.com/scitable/blog/viruses101/are_modified_mosquitoes_the_future

Campbell discusses the world wide spread of dengue and current attempts to prevent the spread.
Campbell, C. "If You're Not Worried About Dengue Fever, Here's Why You Should Be." *Time*. November 18, 2013.

IX. How will students demonstrate understanding?

Successful arrangement of cell images according to directions.
Successfully answering the questions regarding technology and completion of the virtual labs.
Successful completion of the case study.

X. References

Clinic Staff, Mayo. "Dengue Fever." Mayo Clinic. N.p., 19 Sept. 2014. Web. <http://www.mayoclinic.org/diseases-conditions/dengue-fever/basics/definition/con-20032868>.

"Dengue." Centers for Disease Control and Prevention. Centers for Disease Control and Prevention, 26 Mar. 2015. Web. <http://www.cdc.gov/Dengue/>.

"Dengue Guidelines for Diagnosis, Treatment, Prevention and Control." (2009): 1-160. Dengue. World Health Organization, 2009. Web. <http://whqlibdoc.who.int/publications/2009/9789241547871_eng.pdf>.

DeSalle, Rob, and Michael R. Heithaus. *Holt Biology*. Orlando, Fla: Holt, Rhinehart and Winston, 2008. Print

Garcia-Navarro, Lourdes. "Ready, Set, Spray! Brazil Battles Dengue Ahead Of The World Cup." NPR. NPR, 29 May 2014. Web. <http://www.npr.org/blogs/health/2014/05/29/316668643/ready-set-spray-brazil-battles-dengue-ahead-of-the-world-cup>.

Hillis, David M. *Principles of Life*. Sunderland, MA: Sinauer Associates, 2012. Print

Murphy, Kenneth, Paul Travers, Mark Walport, and Charles Janeway. *Janeway's Immunobiology*. New York: Garland Science, 2008. Print.

Reece, Jane B, and Neil A. Campbell. *Biology*. Boston: Pearson Learning Solutions, 2008. Print.

XI. STUDENT SECTION

This section contains:

- Video Reflection Guide
- Immune Cell Images

Video Reflection Guide – Immune System

Name_____ **Block**_____

Respond to the following questions after watching the Immune System Video.
http://www.bozemanscience.com/immune-system

Brief Summary **"In this video presentation I learned…"**

The main points

Innate and Adaptive Cells of the Immune system





What are PCR, RT-PCR and Flow Cytometry?

Continue on to the PCR Virtual lab and answer the following questions.

1) What is the importance of PCR?

2) Where can DNA samples come from for PCR reactions?

3) To start a PCR reaction what do you need to add to the PCR tube besides the DNA?

4) What is the name of the machine that can heat and cool the PCR tube?

5) What happens to DNA at 95^0C?

6) Once the DNA Polymerase locates the primers what does it do?

7) By the end of cycle three, how many copies of the DNA fragment you targeted are there?

III. You now have an understanding of the basic principles of PCR. Read the following short article on Reverse Transcriptase Polymerase Chain Reaction (RT-PCR) and answer the following questions.

http://www.bio.davidson.edu/people/Kabernd/seminar/2002/method/lowry/RTPCR.htm

1) What nucleic acid is detected by this technique?

2) If that nucleic acid is present in the sample being tested, what does it mean for a specific gene that it is associated with and its product?

3) What are the steps in the "RT" portion of RT-PCR?

IV. What is Flow Cytometry?

Read the information on the Flow Cytometry and answer the accompanying questions.
http://www.biotechnologyforums.com/thread-2185.html

1) What is the function of a flow cytometer?

2) What is the basis for the separation of different cell types?

3) Explain two uses for this technique.

Dengue - Is the World Cup Worth the Risk?

Introduction:

October 30, 2007 – Brazil has been elected as the host nation for the 2014 FIFA World Cup. Fast forward to August 20, 2013.

Chris J., an enthusiastic soccer fan, is anxiously awaiting the online ticket sales of the 2014 FIFA World Cup to open up. Chris is a Ph.D. student in the Microbiology and Immunology Department at Upstate Medical University. He spends his summers in Brazil researching the mosquito dengue fever vectors, *Aedes aegypti and Aedes albopictus.* Next summer, while collecting mosquitoes from the City of Natal, he is hoping to see the U.S. play a soccer match in the Arena das Dunas. He will be taking Drew, an undergraduate research assistant, with him to help with collecting the mosquitos and he wants to go to the match as well.

Question: What is dengue?
Answer: Dengue is a vector borne viral disease caused by one of four different serotypes (DENV-1, DENV-2, DENV-3, DENV-4). The virus is a member of the Flaviviridae family.

Question: How is it contracted/transmitted?
Answer: The most common vector is the *Ades aegypti* mosquito, *Ades albopictus* is also a carrier. When an infected mosquito bites a human, the virus can be transmitted through the fluids the mosquito inserts through its proboscis. The mosquito acquires the virus through the blood of an infected human it bites. Human to human transmission is not possible.

Question: Can I get dengue again if I have already been infected once?
Answer: You will have immunity to the specific virus (DENV-1, DENV-2, DENV-3, DENV-4) that you were originally infected with and, if infected again with that serotype, you will not get the disease. However, if you are infected with a different serotype you are at a greater risk of acquiring dengue hemorrhagic fever (DHF). DHF is a more severe form of dengue and can be fatal if medical treatment is not immediately sought.

Question: How can it be prevented?
Answer: As the disease is spread by mosquitoes, eradication of breeding sites is essential. *Ae. aegypti* lays her eggs in containers that hold water (e.g. tires, planters, water drums). However, *Ae. aegypti* are highly adaptable to varying conditions; eggs have been known to survive drought conditions. *Ae. albopictus* prefers cool dark areas such as home closets. The use of air conditioning and screens can prevent adults from finding their human meals.

Question: What are the disease symptoms?
Answer: Many of the symptoms are very similar to malaria or the flu, they include: high fever; severe headache; pain in the joints, muscle, bone or behind the eyes; rash; and mild bleeding. The symptoms of dengue hemorrhagic fever include a fever that lasts from two to seven days with the other symptoms mentioned above. As the fever declines the following symptoms may occur: vomiting, abdominal pain and breathing difficulties. This opens a 24 to 48 hour period when capillaries start to leak fluid. This increase in fluid in the body cavities can lead to the failure of the circulatory system, shock, and death. Skin hemorrhages, along with internal and external bleeding, are also possible.

Question: How is dengue treated?
Answer: There is no specific treatment for the disease. If one suspects they have dengue fever, they can take analgesics, stay hydrated, rest, and speak with their doctor. If they develop more severe symptoms after the fever has subsided, they should go to a hospital. Fluid replacement therapy is appropriate at this point. Currently, there is no vaccine on the market; however, the company Sanofi Pasteur has plans to register the tetravalent vaccine (protects against all 4 dengue strains) in 2015 with a target distribution time frame of late 2015.

Their research went well; they were able to sample adult mosquito populations as well as collecting eggs from a variety of sources. While it was winter in Brazil mosquitoes were still plentiful. On June 16th, the day before they were to leave Natal, they went to the Arena das Dunas to watch the US vs Ghana soccer match. The US won 2-1, what a great way to end their trip!

Symptoms?

Chris and Drew had a non-eventful trip back to Syracuse on the 17th. Chris was feeling a bit "off." He was tired and a bit achy when he got back, but put it down to a hectic few weeks. He went into the lab on the 18th, by mid-day [illegible]

in a day. In the meantime, he should go home making sure that he drinks fluids and, if needed, take acetaminophen.

Chris went to see Dr. Matthew the next day. Dr. Matthew had received the lab reports back. The Real Time RT PCR was negative; it showed no viral RNA in the blood sample. As the virus is only present in the blood for the first five days after infection, this was not an unexpected result. The MAC ELISA, testing for the presence of IgM, was positive for DENV-2. The IgG ELISA was negative for DENV. Dr. Matthew told Chris that it was likely that he has been infected with DENV-2. The negative IgG ELISA result indicates that this probably is a primary not secondary infection. That determination will be made when you come back in 14 days for another blood draw for the second IgG ELISA.

15 days later – Chris now fully recovered from his symptoms returns to Dr. Matthew's office for the results of his second IgG ELISA test. Dr. Matthew tells Chris that his second IgG ELISA was positive. The official diagnosis is "recent probable primary dengue infection."

9) Explain what Chris' final diagnosis is.

The follow up

Dr. Matthew asked Chris if he remembers being bitten by a mosquito. Chris said he couldn't remember being bitten. About 6 days before they left Natal, he and Drew were sampling in an area of the city that had significantly more pools of stagnant water than the other areas that they had sampled. It seemed as if there were clouds of mosquitoes hovering around some of their sampling spots.

Dr. Matthew asked if Chris would be going back to Brazil to gather more samples in the future; reminding him that an encounter with the DENV-1, 3 or 4 serotypes could lead to dengue hemorrhagic fever and potential death. The body's immune system will produce antigens to the specific DENV serotype it was first exposed to but not hr es but

10) If you were Chris would you be concerned about a secondary dengue infection? Why or why not?

EXTENSION

1) Students can read about the dengue virus on the Protein Data Bank web page. Using Jmol a student can manipulate an envelope protein with antibodies bound and the complete Dengue Virus with antibodies attached.
http://www.rcsb.org/pdb/101/motm.do?momID=103

2) Students can construct a paper model of the dengue Virus.
http://www.rcsb.org/pdb/education_discussion/educational_resources/dengue_virus_3Dmodel.pdf

XII. APPENDIX

Virus

Macrophage engulfing a virus

What are PCR, RT-PCR and Flow Cytometry?

TEACHERS ANSWERS

I. The following video from the DNA Learning Center at Cold Spring Harbor, elaborates upon the steps that take place in the typical PCR reaction.

https://www.youtube.com/watch?v=2KoLnIwoZKU

3) To start a PCR reaction what do you need to add to the PCR tube besides the DNA?
(Primer 1, Primer 2, Nucleotides (Adenine, Guanine, Thymine and Cytosine), DNA Polymerase)

4) What is the name of the machine that can heat and cool the PCR tube?
(Thermal Cycler)

5) What happens to DNA at 95^0C?
(It denatures.)

6) Once the DNA Polymerase locates the primers what does it do?
(Add complementary nucleotides to the single stranded DNA.)

7) By the end of cycle three how many copies of the DNA fragment you targeted are there?
(Two)

III. You now have an understanding of the basic principles of PCR. Read the following short article on Reverse Transcriptase Polymerase Chain Reaction (RT-PCR) and answer the following questions.

http://www.bio.davidson.edu/people/Kabernd/seminar/2002/method/lowry/RTPCR.htm

1) This technique detects the presence of what nucleic acid?
(mRNA or messenger RNA)

2) If that nucleic acid is present in the sample being tested what does it mean for a specific gene that it is associated with and its product?
(That gene is being expressed and therefore whatever the gene codes for (usually a protein) will be made.)

3) What are the steps in the "RT" portion of RT-PCR?
(Use reverse transcriptase and the appropriate primer to locate the desired mRNA sequence, this will transcribe a complimentary copy of DNA (cDNA), using standard PCR techniques the DNA is replicated using primers and Taq Polymerase.)

IV. What is Flow Cytometry?

Read the information on the Flow Cytometry and answer the accompanying questions.

http://www.biotechnologyforums.com/thread-2185.html

1) What is the function of a flow cytometer?
(To separate out specific cell types.)

2) What is the basis for the separation of different cell types?
(Their size, shape and color that they fluoresce.)

3) Explain two uses for this technique.
(

Detection of the Prevalence of Lyme Disease in Local Canine Populations
Teacher Section

Introduction

What is Lyme d

distinguish between animals that have had prior exposure from vaccinated or non-infected animals. Another antigen used in ELISAs for the detection of Lyme infection is Decorin binding protein A (DbpA). DbpA is a bacterial outer membrane lipoprotein that facilitates the attachment of *B. burgdorferi* to the tissue of its host. IgG activity is detectable throughout the progression of the disease. In contrast to the C_6 peptide, both vaccinated and infected animals produce DbpA antibodies, and the DbpA ELISA, by itself, cannot distinguish between infected and vaccinated animals.

Canines and Lyme Disease

The studies in this project will identify antibodies (IgG) in the serum of local dogs. Dogs have been chosen for these studies as Lyme disease is one of their most common arthropod borne diseases. Central New York has seen a steady rise in the number of infected dogs over the past several years. This rise has also been mirrored in the human population. Detection of Lyme disease in canines can act as a surrogate detection assay for the risk of human exposure to the disease as both dogs and humans inhabit and exercise in the same areas.

Objectives

Your local veterinary clinic routinely collects blood samples from its canine patients. Utilizing the technique of ELISA, you will determine if canines carry antibodies to the Lyme pathogen in their serum. If so, these studies will demonstrate that the dogs have been exposed to the Lyme pathogen.

Advanced Preparation

How to Make Buffers

1) Phosphate-Buffered Saline (PBS- 1X) 1 Liter

NaCl	8 g
KCl	0.2 g
Na_2HPO_4	1.44 g
KH_2PO_4	0.24 g

Dissolve the reagents in 800 ml of distilled H_2O and adjust the pH to 7.4 with HCL. Add additional H_2O to make 1 Liter.

2)

Materials List

Microtiter Plates
ELISA Wash Buffer
ELISA Blocking Solution
10% Sodium Azide (NaN_3) in milli-q water
Capture antibody - DbpA protein
Canine Serum or Plasma
Goat anti-canine IgG (H+L)-AP
AP Substrate
Plastic wrap
Eppendorf tubes
Pipettors and tips: P1000, P200, P20
Bottles to hold buffers
Sterile pipettes 1ml, 5 ml
Elisa Plate Reader

Materials Price List

Product	Supplier	Order #	Quantity	Price w/o shipping
Phosphatase (AP) Substrate	Sigma Aldrich	S0942-200TAB	1 bottle	$112.00
Tween 20	Sigma Aldrich	P1379-25ml	25 ml	$11.40
Sodium Azide (NaN_3)	Sigma Aldrich	S200205G	5 gm	$12.50
Glycine	Sigma Aldrich	410225-50G	50 gm	$35.10
$MgCl_2$	Sigma Aldrich	M8266-100G	100 gm	$52.10
$ZnCl_2$	Sigma Aldrich	229997-10G	10 gm	$113.50
NaCl	Sigma Aldrich	S7653-250G	250 gm	$35.60
KCl	Sigma Aldrich	P9333-500G	500 gm	$79.90
				$37.90

ELISA Protocol

Prior to starting the experiment, label a mock ELISA plate sheet (see end of protocol).

Day 1 (Coating of the Plate)

1) Coat wells of the microtiter plate with 100ul of a 1μg/100μl capture antibody solution.
2) Wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 2 (Blocking of the Plate)

3) Wash plate with ELISA wash buffer 3 times by filling each well with 100 μl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
4) Block wells by filling each well with 100 μl Blocking Buffer and incubating for 1 hour at room temperature.
5) Wash plate with ELISA wash buffer 2 times by filling each well with 100 μl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
6) Place 100 μl ELISA wash Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 3 (Addition of the Primary Antibody – the canine serum/plasma)

7) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
8) For each different canine serum/plasma sample follow the following dilution scheme

In well A pipet 196 μl ELISA wash buffer and 4 μl of serum/plasma.
In wells B-D pipet 100 μl ELISA wash buffer.
Take 100 μl of liquid from well A and add it to well B, pipet up and down twice to ensure mixing.
Take 100 μl of liquid from well B and add it to well C, pipet up and down twice to ensure mixing.
Take 100 μl of liquid from well C and add it to well D, pipet up and down twice to ensure mixing.



9) Cover plate and Incubate at room temperature for 1 hour.
10) Wash plate with ELISA wash buffer 2 times by filling each well with 100 µl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
11) Place 100 µl Blocking Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 4 (Addition of the Secondary Antibody)
11) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
12) To the wells add 100 µl of enzyme coupled AP secondary antibody (goat-anti-canine-IgG-AP) diluted 1:1000 in ELISA Wash Buffer. Incubate at room temperature for 1 hour.
13) Wash plate with ELISA wash buffer 2 times by filling each well with 100 µl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
14) Place 100 µl ELISA Wash Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 5 (Addition of the Substrate)
15) Prepare AP substrate; add one pellet of Alkaline Phosphatase to 3 mls of Glycine Buffer. This is enough for approximately 30 wells.
15) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
16) Add 100 µl of AP substrate to each well.
17) Allow to develop 20 minutes and immediately read on an ELISA plate reader or quench reaction with 50 µ

Samples that have curves with significantly higher OD values than that of the control will be considered to be positive for Lyme disease.



Format for Laboratory Report

Students will use the mini-poster format. The board of the poster consists of two manila folders that overlap giving the author three surfaces to affix their text. This method is adapted from A Handbook of Biological Investigation. Harrison W. Ambrose III and Katharine Peckham Ambrose. 1995, Hunter Textbooks and Brad Williamson.

Headings:

Title and Author. The title should be descriptive and include the variables that are manipulated.

Abstract. The abstract summarizes the experiment including background, results and conclusions. It should be one to two paragraphs in length.

Introduction. The introduction includes background on the topic and the hypothesis.

Methodology. This is in paragraph form and includes enough information that a reader familiar with the field could replicate the experiment and data analysis.

Results. Use graphs, tables and charts to help clarify the data. Include a short summary of what the data mean under the visual used.

Discussion. Here you discuss what your results mean in relationship to your hypothesis. Point out the significance of your findings. If the results are not what was expected, attempt to explain why, and point out other research avenues.
Literature Cited. Include all published works mentioned in your report.

References

Arnaboldi, P. M., M. Sambir, and R. J. Dattwyler. "Decorin Binding Proteins A and B in the Serodiagnosis of Lyme Disease in North America." Clinical and Vaccine Immunology 21.10 (2014): 1426-436. Web. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4266351/>.

"Http://bakerinstitute.vet.cornell.edu/animalhealth/page.php?id=1101." Baker Institute/Animal Health. Cornell University College of Veterinary Medicine, 2007. Web.

Levy, S. A., T. P. O'connor, J. L. Hanscom, P. Shields, L. Lorentzen, and A. A. Dimarco. "Quantitative Measurement of C6 Antibody following Antibiotic Treatment of Borrelia Burgdorferi Antibody-Positive Nonclinical Dogs." Clinical and Vaccine Immunology 15.1 (2008): 115-19. Web. https://www.idexx.com/pdf/en_us/smallanimal/snap/lyme/quant-c6-paper.pdf

Detection of the Prevalence of Lyme Disease in Local Canine Populations
Student Section

Introduction

What is Lyme disease and what are its symptoms?

Lyme disease is the most common arthropod borne disease in humans. It is highly prevalent in the canine population, particularly in the Northeastern US. The disease is caused by the bacterium *Borrelia burgdorferi*, which resides in the midgut of the deer tick. When a tick bites its host (i.e. takes a blood meal) *B. burgdorferi* moves into the salivary glands of the tick and then, during tick feeding, *B. burgdorferi* moves into its host. The likelihood of infection increases with the duration of tick feeding. It is thought that before 24-48 hours of feeding there is little risk of infection.

What is the Life Cycle of a Deer Tick?

Deer ticks have a life cycle that spans two years, three hosts, and four developmental stages. Eggs are laid in the spring, but the larvae that develop do not carry enough bacteria to cause an infection. Larva in the Northeastern United States feed primarily on the white footed mouse which can harbor the bacteria for long periods before exhibiting symptoms of the disease. The larvae become infected with the bacteria by feeding on the blood of the infected mouse. As winter approaches the larvae leave the mice and enter a resting phase.

The following spring the larvae molt into nymphs. During the spring and summer months the nymphs feed on new hosts, including mice, humans, and dogs. A nymph, during its four day feeding period can be infected by the host it is feeding on, if the host is already infected with *B. burgdorferi*. The nymph, if previously infected in the larval stage, can also infect its host.

During the fall the nymphs molt again, into adult ticks. On average 50% of adult ticks in infected areas in the Northeast US carry the infection. The adult ticks, which can commonly be found on shrubs and other vegetation, will attach to deer and other large animals.

Adult ticks will mate on their new host. Males tend to stay on the host and die whereas females feed for five to seven days then drop off into the leaf debris, where they over winter. The following spring the females will lay eggs, completing the life cycle.

How is Lyme Disease Diagnosed?

distinguish between animals that have had prior exposure from vaccinated or non-infected animals. Another antigen used in ELISAs for the detection of Lyme infection is Decorin binding protein A (DbpA). DbpA is a bacterial outer membrane lipoprotein that facilitates the attachment of *B. burgdorferi* to the tissue of its host. IgG activity is detectable throughout the progression of the disease. In contrast to the C_6 peptide, both vaccinated and infected animals produce DbpA antibodies, and the DbpA ELISA, by itself, cannot distinguish between infected and vaccinated animals.

Canines and Lyme disease

The studies in this project will identify antibodies (IgG) in the serum of local dogs. Dogs have been chosen for these studies as Lyme disease is one of their most common arthropod borne diseases. Central New York has seen a steady rise in the number of infected dogs over the past several years. This rise has also been mirrored in the human population. Detection of Lyme disease in canines can act as a surrogate detection assay for the risk of human exposure to the disease as both dogs and humans inhabit and exercise in the same areas.

Objectives

Your local veterinary clinic routinely collects blood samples from its canine patients. Utilizing the technique of ELISA, you will determine if canines carry antibodies to the Lyme pathogen in their serum.

Materials List

Microtiter Plates
ELISA Wash Buffer
ELISA Blocking Solution
10% Sodium Azide (NaN_3) in milli-q water
Capture antibody - DbpA protein
Canine Serum or Plasma
Goat anti-canine IgG (H+L)-AP
AP Substrate
Plastic wrap
Eppendorf tubes
Pipettors and tips: P1000, P200, P20
Bottles to hold buffers
Sterile pipettes 1ml, 5 ml
Elisa Plate Reader

ELISA Protocol

Prior to starting the experiment, label a mock ELISA plate sheet (see end of protocol).

Day 1 (Coating of the Plate)

1) Coat wells of the microtiter plate with 100ul of a 1μg/100μl capture antibody solution.
2) Wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 2 (Blocking of the Plate)

3) Wash plate with ELISA wash buffer 3 times by filling each well with 100 μl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
4) Block wells by filling each well with 100 μl Blocking Buffer and incubating for 1 hour at room temperature.
5) Wash plate with ELISA wash buffer 2 times by filling each well with 100 μl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
6) Place 100 μl ELISA wash Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 3 (Addition of the Primary Antibody – the canine serum/plasma)

7) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
8) For each different canine serum/plasma sample follow the following dilution scheme

In well A pipet 196 μl ELISA wash buffer and 4 μl of serum/plasma and.
In wells B-D pipet 100 μl ELISA wash buffer.
Take 100 μl of liquid from well A and add it to well B, pipet up and down twice to ensure mixing.
Take 100 μl of liquid from well B and add it to well C, pipet up and down twice to ensure mixing.
Take 100 μl of liquid from well C and add it to well D, pipet up and down twice to ensure mixing.

9) Cover plate and Incubate at room temperature for 1 hour.
10) Wash plate with ELISA wash buffer 2 times by filling each well with 100 µl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
11) Place 100 µl Blocking Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 4 (Addition of the Secondary Antibody)
11) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
12) To the wells add 100 µl of enzyme coupled AP secondary antibody (goat-anti-canine-IgG-AP) diluted 1:1000 in ELISA Wash Buffer. Incubate at room temperature for 1 hour.
13) Wash plate with ELISA wash buffer 2 times by filling each well with 100 µl buffer, flicking plate face down into the sink and then tapping gently on several paper towels to remove residual fluid.
14) Place 100 µl ELISA Wash Buffer in wells and wrap plate in plastic wrap and incubate overnight at 4^0C.

Day 5 (Addition of the Substrate)
15) Prepare AP substrate; add one pellet of Alkaline Phosphatase to 3 mls of Glycine Buffer. This is enough for approximately 30 wells.
15) Flick plate face down into the sink and then tap gently on several paper towels to remove residual fluid.
16) Add 100 µl of AP substrate to each well.



Format for Laboratory Report

You will use the mini-poster format. The board of the poster consists of two manila folders that overlap giving the author three surfaces to affix their text. This method is adapted from A Handbook of Biological Investigation, Harrison W. Ambrose III and Katharine Peckham Ambrose. 1995, Hunter Textbooks and Brad Williamson.

Headings:

Title and Author. The title should be descriptive and include the variables that are manipulated.

Abstract. The abstract summarizes the experiment including background, results and conclusions. It should be one to two paragraphs in length

Introduction. The introduction includes background on the topic and the hypothesis.

Methodology. This is in paragraph form and includes enough information that a reader familiar with the field could replicate the experiment and data analysis.

Results. Use graphs, tables and charts to help clarify the data. Include a short summary of what the data mean under the visual used.

Discussion. Here you discuss what your results mean in relationship to your hypothesis. Point out the significance of your findings. If the results are not what was expected then attempt to explain why and point out other research avenues.

Literature Cited. Include all published works mentioned in your report.

References

Arnaboldi, P. M., M. Sambir, and R. J. Dattwyler. "Decorin Binding Proteins A and B in the Serodiagnosis of Lyme Disease in North America." Clinical and Vaccine Immunology 21.10 (2014): 1426-436. Web. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4266351/>.

"Http://bakerinstitute.vet.cornell.edu/animalhealth/page.php?id=1101." Baker Institute/Animal Health. Cornell University College of Veterinary Medicine, 2007. Web.

Levy, S. A., T. P. O'connor, J. L. Hanscom, P. Shields, L. Lorentzen, and A. A. Dimarco. "Quantitative Measurement of C6 Antibody following Antibiotic Treatment of Borrelia Burgdorferi Antibody-Positive Nonclinical Dogs." Clinical and Vaccine Immunology 15.1 (2008): 115-19. Web. https://www.idexx.com/pdf/en_us/smallanimal/snap/lyme/quant-c6-paper.pdf