Using the “Smurf Transition” to Study Leaky Gut in a Fruit Fly Model

Gregory Shenk
Capitol Region Education Council – Academy of Aerospace and Engineering
1101 Kennedy Rd
Windsor, CT 06095
gshenk@crec.org

Mentors: Kamal Khanna, Department of Immunology
University of Connecticut Health Center
Pablo Romagnoli
Instituto Universitario de Ciencias Biomédicas de Córdoba, Argentina

With funding from The American Association of Immunologists

Table of Contents

Science Background 2
Student Outcomes 4
Course Placement 4
AP Biology Curriculum Standards 4
AP Biology Science Practices 5
Time Requirements 5
Advance Preparation 5
Materials and Equipment 6
Student Prior Knowledge and Misconceptions 7
Daily Unit Plans 8
Student Section 10
    Rationale 10

model organisms to test their hypotheses about what goes wrong in inflamed or “leaky” guts and to discover ways that these conditions can be avoided or treated with fewer complications.

The workhorse in this research is the mouse and many variants that mimic human genetic vulnerabilities in IBDs are now available; some even express human proteins so researchers can better understand why certain heritable gene variants make patients more susceptible to IBD. Not surprisingly, the malfunctioning proteins are often found to play a role in the maintenance of the tight junctions, production of epithelial secretions (mucus, the complex carbohydrates that support our friends when our diet cannot, and antimicrobial peptides). They may also be the signals or receptors that our epithelial and immune cells use to monitor and regulate our microbiome. These weak links become therapeutic targets. Unfortunately, experimenting with mice is not possible in most high schools so this unit plan was written to utilize fruit flies as a model organism. Fruit flies harbor a diverse assemblage of microbes (albeit less diverse than vertebrates) in a gut with a single-layer of epithelial cells that is continually renewed and that interacts dynamically with the microbiome using some of the same signaling complexes our cells use.[8] The use of fruit flies as a model organism for IBD research is relatively (e as)h

Velasquez-Manoff, M. 2015. Gut microbiome: the peacekeepers. Nature. 518:S3-S11.

Abreu, M.T. 2010. Toll-like receptor signaling in the intestinal epithelium: how bacterial recognition shapes intestinal function. Nat Rev Immunol. 10:131-44.

Philpott, D.J., M.T. Sorbara, S.J. Robertson, K. Croitoru and S.E. Girardin. 2014. NOD proteins: regulators of inflammation in health and disease. Nature Reviews. 14: 9-23.

Hong, C. and P. Tontonoz. 2008. Coordination of inflammation and metabolism by PPAR and LXR nuclear receptors. Current Opinions in Genetics and Development. 18:461-467.

Materials and Equipment

The materials I used in developing this unit plan were either standard resources the teacher will probably already have (e.g. dissecting microscopes) or are inexpensive. Many pet owners raise their own fruit flies to feed their pets so it's relatively easy to find simple instructions for inexpensive fruit fly feeding media, and your local pet store will probably be able to supply flightless fruit flies. The dye used to detect a leaking gut is actually food coloring and the concentration of that obtained in a grocery store will work just fine. Also, DSS is the standard chemical used to induce inflammation in the guts of laboratory animals, but I recommend dish soap as it will be mu

Daily Unit Plans

2. When finished with discussion of the paper, students should incorporate new information into their models.
3. Students watch video on NOD-like receptors in class: (https://www.youtube.com/watch?v=HSk90Yx4ulE)
4. Students take home NOD-like receptor and Nuclear Receptor papers and do their best to determine what they're about from looking at and reading the captions for the images. The Butyrate in the "Your Microbes at Work" image from paper given on Day 1 is an example of a nuclear receptor ligand.

Day 5

1. Class begins with discussion of the role of NOD and NRs in immunity.
2. Students conduct internet research to identify factors which may either increase or decrease gut inflammatory response and propose a new experiment to test effects of their factor on speed of Smurf Transition.

Day 6

1. Teacher introduces cell biologist's toolkit.
2. Students begin preparing three minute talks on how they might be able to use these tools to study the effects of their treatments.

Day 7

1. As summative assessment, students present their three minute talks, field questions, and respond to feedback.

Student Section

Rationale

You may already be somewhat familiar with the important job your immune system does in helping you avoid getting sick because you had to show proof of vaccination in order to enroll in school, and you likely get vaccinated every year against the seasonal "bugs." But did you know that your gut is the most important site of defense in your whole body? To understand why this is, consider the fact that the thin layer of cells lining your gut has to provide passage for the molecules you absorb from your food and therefore *must be* more permeable than other epithelial barriers. This greater permeability to food molecules presents a potential passageway for pathogens to enter the body and it is for this reason that natural selection produced such a formidable defensive arsenal along the length of our intestines.

Unfortunately, many people around the globe suffer from excessive activation of these defenses in the presence of low-level threats or even in the complete absence of threats, and they may experience the unpleasant effects of their body's response in the form of Inflammatory Bowel Disease (IBD). Consequently, many researchers are engaged in the work of trying to work out the cellular details of these inflammatory responses so that one day those who suffer from them may be able to enjoy relief from treatments aimed at better controlling the inflammatory pathways.

Since most experiments on humans are considered unethical, researchers who want to better understand these and other disorders impacting human health use other organisms as "models." The mouse is one of the most commonly used models because it's a mammal, and its responses to experimental manipulation are more likely to resemble a human's responses to the same treatments since we share a recent ancestor. However, working with mice at the high school level is almost impossible, so you will use a more distant relative which doesn't currently enjoy the protections that mice do. Your model for studying gut immunology will be the lowly fruit fly, *Drosophila* sp.

Researchers trying to understand what happens in our bodies at the end of life have been using fruit flies as models for many years. While searching for ways to study the cellular events that precede death, they discovered that the gut becomes inflamed and breaks down hours before the fly actually dies and before the fly shows an obvious decline in activity or vigor. To help them identify which flies were undergoing this transition, they found that food dyes could be added to a fly's food without harming the flies, and its leakage into the abdominal cavity could be easily observed under a dissecting microscope. They began using the FD&C blue dye used in coloring food as their marker and dubbed the transition from an intact to a leaky gut the "smurf transition" assay. You'll be using this assay to determine which of your treatments trigger gut inflammation or "leaky gut."

Before you begin this unit, you will watch some videos introducing the lymphatic system and the immune system, and you will answer questions as you watch. This is meant to give you a quick introduction to the function and complexity of the immune system. You will then use a

microscope to look at two fruit flies that have been feeding on a growth medium to which blue food coloring was added. One was also fed in a medium to which a detergent was added. You should be able to see a difference between them. One will have the food coloring localized in its digestive tract and the other will have blue diffused throughout its abdomen.
Through a series of videos, images, and papers that your teacher will share, you and your classmates will develop a working model of inflammatory bowel disorder that can explain these conditions at the level of individual cells. Your models will reflect the array of different possible environmental triggers for inflammation, the receptors and pathways activated by them, and the resulting change in phenotype at the tissue level. After developing this model, you will have a chance to set up your own experiment using fresh fruit flies and propose a more advanced analysis using modern molecular tools to reveal just exactly what may be going on at the level of gene activation within distinct cell populations.

Day 1: Building a Model of Leaky Gut in a Fruit Fly

Smurf Flies! Pair up and obtain each of the two fruit flies that your teacher has prepared for you (one fed on a standard medium with food color added and one fed on a medium with food color and detergent). Quickly inspect both fruit flies using a stereomicroscope (or magnifying glass if stereomicroscopes are not available). Write down your observations. What do you see? (Don't worry about trying to explain why. That's where we're going next!)

What's a "leaky gut?" Now your teacher will ask you to view a brief informational video about Inflammatory Bowel Disease or IBD (

Day 2: The Role of Immune Cell Signaling Mechanisms in IBD

A more detailed look at the defenses of the gut epithelium: Watch the video "Immunology of the Gut Mucosa" (https://www.youtube.com/watch?v=gnZEge78_78) and add details about the dynamic role of the gut mucosal immune system to your model of the gut epithelium. As your previous paper may be getting pretty crowded and messy, you may ask your teacher for a fresh piece of poster paper if necessary.

Walk around! At this point, your group has been working in isolation from other groups to come up with the most comprehensive and up-to-date model of the gut epithelium and your ideas are limited to what the members of your group saw and remembered from the videos, pictures and articles they saw. But are there things you missed? Might other students have seen things or thought for things that your group members didn't? This is a good time to find out! Get up, take your notebook, and walk around to look at the work that has been done by other groups. Write down anything you see that your group didn't include but probably should have. When you get back to your own group's poster, add it in!

Talk about it! Scientists develop their understanding of the natural phenomena they study in much the same way you are – by trying to come up with an explanation for what they see, gathering more information and improving their understanding. But an important part of this process is sharing their discoveries and insights with other scientists and studying the work of other scientists to find out what their colleagues have learned in their own investigations. This exchange often takes place at professional meetings where they can watch each other give presentations and talk to each other while looking at posters that summarize their results. You'll try to emulate this now by discussing your models as a whole class.

Wait! There's more! So far you should have a pretty good idea of the way different cells are arranged within and adjacent to the gut epithelium as well as what the different cell types here do. But you should now be wondering about some of the specific mechanisms involved. For instance, if one immune cell sends a signal and the cell receiving it undergoes a phenotypic change or produces a product of its own in response, how is this happening? Your teacher will now share a video with you about the Toll-Like Receptors or TLRs (https://www.youtube.com/watch?v=GXECgTLGLtI) and for homework you will be assigned to become very familiar with *one portion* of a scientific paper written to review what was known about the function of TLRs at the time the paper was written. To help you, the teacher will provide a reading guide and a glossary for technical terms and acronyms you're unlikely to know. Do the best you can because you'll have to teach your classmates what you know the next time you meet!

We're smarter together: So far you've been adding to your understanding of the gut epithelium somewhat like a scientist might do. The model you're holding in your head has become increasingly sophisticated as you've added details to your understanding. The paper you are being given at the end of this class was full of details; too full, in fact, for you to really

is a collaborative process and, when scientists collaborate, they can achieve far more than they ever could have done working in isolation. Because your teacher assigned you to read just one part of the paper and you will become the expert on a certain piece of the TLR response puzzle, your task in the next period will be to show other class members what you have learned and they will do the same.

Days 3&4: Reading a Scientific Paper

Time to put on your teacher-scientist hat! Since visualization helps us all understand things better, you are asked to add your knowledge to a classroom model of the gut mucosal immune system. Take your time. This may take a couple class periods depending on how much you and your classmates have to say! When the class has finished fleshing out the details of Toll-Like Receptor signaling, each group should meet for a few minutes to discuss their models and try to reach a consensus on how this new knowledge may be incorporated into these models. For the next class, students will discuss two other classes of receptors that may play a role in the cellular response to pathogen-derived factors. One of these is a class of intracellular receptors called NOD-like receptors. You'll watch the video about them in class before you leave (https://www.youtube.com/watch?v=HSk90Yx4ulE) and skim through the scientific review paper your teacher gives you about them to get a handle on how they compare to TLRs. This paper is like the one you just read about TLRs but see if you can get the "gist" by flipping through the paper and studying the diagrams. Do the same with the paper on nuclear receptors. These are yet another class of receptor that may enable your cells to recognize and respond to bacteria that are nearby. They're the receptors used by the immune cells to detect the short chain fatty acid metabolites (e.g. butyrate) shown in 1(r)-6(s)-2(d6(e)11(c11(k( )11(t)l(he on)11(n)11

Transition" technique and variables of your own choosing. As you saw at the beginning of this unit, detergent added to the fly's food will disrupt the epithelium and induce a leaky gut. But what else can make the gut leaky? Are there things that help protect it? To get some ideas, you might begin by doing an internet search for foods people eat to reduce inflammation. Thinking about your model and the different

Appendix

Crash Course Video Homework: The Lymphatic and Immune Systems

Name:_____Period:_____

Date:_____

Lymphatic System (https://www.youtube.com/watch?v=I7orwMqTQ5I)

1. What are the primary roles of the lymphatic system?

2. What are the three main parts of the lymphatic system?

3. What do the lymph nodes and lymphatic glands do?

4. How does lymph move through the lymphatic system?

5. What can happen in a lymph node if an invader arrives there?

6. What are MALTS, where are they found and what do they do?

7. What function does the appendix have?

Immune System part 1 (https://www.youtube.com/watch?v=GIJK3dwCWCw)

1. How does the immune system differ from other organ systems?

2. What is the first line of defense in your immune system?

3. What are some of the innate defenses you have in your mucosal tissues?

4. What's the 2nd line of innate defense, what are some of the cells it has and what can they do?

5. Why is MHC 1 important in innate immunity?

6. What's the inflammatory response? What are the jobs of the temperature changes and histamines?

7. What will cause cells to release pyrogens and what do they do?

8. What if the pyrogenic response doesn't do the job?

Immune System part 2 (https://www.youtube.com/watch?v=2DFN4IBZ3rI)

1. What makes the adaptive or acquired immune system different than the innate defenses?

2. What are the two components of the adaptive immune system and what do they do?

3. How do vaccinations work?

4. Where do the B-cells mature and what does this entail? What do they produce on their surface and where do they go after they mature?

5. What activates the B-cells and what will they do as a result?

6. How do plasma cells and memory cells differ in terms of function?

7. Antibodies can't directly kill invaders but what can they do?

8. What's the secondary immune response and how do vaccines activate it?

9. Why do we need new vaccinations to be protected from some pathogens?

10. What is passive immunity and how does it work?

Reading Guide for “Gut Microbiome: The Peacekeepers”
Name: _____Period:_____ Date:_____

1.

8.

16. What did researchers find when they studied patients with and without inflammatory bowel disease who possessed the same NOD2 mutation that is frequently associated with the disease?

17. What does the fact that not all of these individuals had inflammatory bowel combined with the much lower incidence of the disease in adults who grew up on farms suggest about the way we may want to think about managing this disease in the future?

18. Throughout this article we've seen references to ecological concepts and one of the most talked about concepts among modern ecologists is the importance of high species diversity in maintaining robust, stable ecosystems. What do the comparisons of gut microbiomes in North Americans to those of rural residents in Africa and South America imply about the bacterial ecosystems of our guts with respect to this species richness-ecosystem resilience principle?

19. What are the products of bacterial fermentation of soluble fiber in our guts that are thought to contribute to a more stable immune phenotype and how do scientists think this may work? What else does Mazmanian think may be at play?

20. What do studies of mice and people show with respect to the challenge of restoring a healthy microbiota with changes in diet?

21. What did researchers observe when they transplanted microbes from healthy donors into patients with metabolic syndrome?

22. Why is Sonnenburg considering a strategy of exterminating the existing microbiome in some patients, administering immune suppressants and then introducing a favorable microbiome? What risks might this entail?

23. How does McFall-Gnai think animal gut evolution may relate to competition in an environment of scarcity and how did the symbiosis support a mobile lifestyle if she's right?

Reading guide for: Toll-like receptor signaling in the intestinal epithelium: how bacterial recognition shapes intestinal function
Name: _____**Period:**_____
Date:_____

Directions: Your teacher will assign you to read one or more sections of this paper. You must understand it well enough to explain it to your peers. There will be some terms and acronyms you may not be familiar with so a glossary has been created to help you as you go. You will also want to make sure you understand how the figures in the paper illustrate the concepts, structures and processes you're reading about because you'll need to use these figures to teach your part of the paper to your classmates. You may find it helpful to redraw the figures forr ind itf -6(t)bk57n?(g)

G4(t)od 1(t)-en-6(oc)-2(e)9(e)maou geadgigors and

9. What's the difference between commensal and pathogenic flora?

10. What effects are TLRs known to have in cells *in general* that the author says are crucial aspects of gut homeostasis?

11. So in addition to their role in maintaining a healthy gut, TLR signals appear to pose hazards through chronic inflammation like we're learning about in IBD pathology. What other disease state is the author interested in learning more about treating as a result of an improved understanding of TLR signaling and gut inflammation?

Sensing bacteria by the intestinal epithelium

12. Pause for a moment to examine figure 2 before you begin reading this section. Look at all the different types of TLRs found in the gut epithelial cells and the differences in the way they're distributed. What do you think the significance of this diversity may be?

13. What does the author mean when she says the IECs are functionally and structurally polarized and how does she claim that this polarity is maintained?

14. What function did this polarity in structure combined with the mucus layer on the epithelium evolve to support and what evidence does the author cite to point out that it "leaks" in healthy individuals but even more so in people with non-functioning TLRs?

15. What would detection of PAMPs by basolateral TLRs be an indication of?

16. The function most biologists from your teacher's generation or before would remember learning that TLRs are critical for detecting invaders and warning the immune system so it could mount an appropriate response. What would these biologists probably find surprising about the expanded role of TLRs in the gut lumen?

TLR expression in the intestine

17. What did scientists set out to find out with regard to commensal bacteria and TLR expression in IELs?

18. What techniques did they have to use and why?

19. What did they find?

20. Sometimes cell biologists will measure gene expression both in terms of the relative amount of mRNA expression and the relative amount of the protein or polypeptide corresponding to the mRNA that may be present in the cell. Why measure both?

21. Aside from determining how abundantly a gene's expression is in terms of the amount of mRNA or protein, what else would a biologist want to know about the gene's product(s)?

22. How does having an inflammatory bowel condition appear to affect TLR expression?

23. Cytokines are often thought of as pro-inflammatory and TLRs as being "for" alerting our immune systems that pathogens are about. But what alternative role may some of the cytokines and some of the TLRs have in gut homeostasis?

24. Which cytokines and which TLRs may be helping keep the calm in healthy guts?

25. What are germ-free mice and what did studies comparing their IECs to those of guts with conventional flora show?

26. What kind of information can investigators get from using immunohistochemistry instead of simply studying gut scrapings?

<u>TLR expression by cell lineage in the intestine</u>

27. So far we've seen that there are multiple TLRs expressed in IECs and their diversity in kind may reflect a similar diversity in functions, with some promoting and some s1(uTj -92 Tc s)-2rS(t)Tw -1.9.002 0

Sensing intestinal injury through TLRs

52. Where on the intestinal epithelium are the crypts and how do the cells here give rise to the rest of the epithelial cells? What are WNT, Notch and Hedgehog factors and what role might they play in this differentiation?

53. What are the components of the stem cell niche that lie beneath the crypts?

54. How do germ-free mice compare to those with a conventional flora in terms of the rate of cell proliferation in the crypts and of what interest are the TLRs with respect to this difference?

55. What did investigators hope studying mice that either lacked functional TLRs or MyD88 would show them?

56. How does proliferation and barrier integrity in mice with no TLR signaling compare to normal mice under normal (non-injury and in the presence of conventional flora) conditions?

57. Why, according to the author, might this be the case in light of other evidence that IECs sense bacteria through this signaling system?

58. What other signaling system does she think might explain this?

59. What is DSS used for in gut epithelia experiments and what does treatment of mice with deficiencies in T

62. TLR4 activation of IECs also appears to lead to recruitment of macrophages to the injury site and mesenchymal stromal cells to the stem crypt. What role do they appear to play in repair when they arrive and what molecules seem to be needed both for their relocation to occur and to trigger their secretion of proliferation-inducing factors?

63. What was the design of the experiments that were used to show that COX2 and MyD88 were critical to inducing the signals for cell proliferation response in injury conditions?

64. What are the ligands for TLR2 and 3 and how does their activation appear to modulate epithelial injury?

65. What ligand have researchers found to induce protection of the epithelium through the TLR9 receptor, which cells are activated by this binding and how do they produce their protective effects?

<u>Radiation induced injury</u>

66. Patients who are undergoing radiation treatment for tumors often experience hair loss, compromised immunity and gut irritation due to the toxicity of the treatments to specific cell populations and their therapy must be suspended while these tissues recover. What makes all of these cells so susceptible to radiation damage?

67. How might the TLR signaling research that was reviewed in this paper (and follow up studies) be applied to helping physicians lengthen the amount of time patients can endure chemotherapy treatments without gut injury?

68. How did LPS treatment of mice prior to radiation treatment protect them?

69. What effect did administration of flagellin prior to radiation have in mice and monkeys and what are the components of the signaling pathway?

70. In addition to direct oral or systemic administration of PAMPs, how might physicians eventually be able to activate protection of the gut in radiation patients?

Necrotizing enterocolitis

71. What are the effects of NEC in infants and, according to studies in mice, what role do TLR2, TLR4 and TLR9 appear to play?

72. With respect to NEC prevention and treatment, what kinds of protocols appear promising as a result of increasing knowledge about TLR signaling and its role in both the injury and protection of the gut epithelium?

TLRs regulate barrier function

73. What are tight junctions composed of and what is their function?

74. What effect did the introduction of a commensal bacterial species have on the expression of genes for these tight junctions in the gut epithelia of germ-free micec 0 Tw-87.6b924j 0.272 0 Td ( )Tj EMC /P <

81. What are the PAMPs th

91. How can experiments that involve localized deletion of NFκB or TAK1 functioning in IECs indicate the importance of TLR activity in barrier function?

92. If bacteria can induce IECs to produce the cytokines TSLP, TGF-beta and IL-25 and each of these in turn has the ability to modulate the level of response by the anti-pathogenic helper T cells, T_H2 and T_H17, and decrease dendritic cell vigilance, what kind of distinction must these IECs also be helping the immune system to make among the species in the gut lumen?

Trefoil factor

93. How do the cells on the margins of an epithelial lesion change to promote healing (restitution) and what role does TFF3 in the mucus layer that the goblet cells produce seem to play

99. How do cytotoxic T cells and NK cells attack tumor cells?

100. Why should the final paragraph give pause to anyone thinking that nonspecific downregulation or upregulation of TLR activity would be a promising approach to treating colon cancer? What might be a better approach given what is known so far?

Glossary for: Toll-like receptor signaling in the intestinal epithelium: how bacterial recognition shapes intestinal function

Absorptive enterocytes – IECs that make up the majority of the intestinal epithelium and play a primary role in nutrient absorption

Amphiregulin – an EGF that promotes proliferation of gut epithelial cells

APRIL – a proliferation-inducing ligand; a cytokine from the TNF superfamily; a B-cell stimulating factor that promotes B-cell survival and class switching

Basement membrane – an extracellular mesh made up of protein fibers which provides a scaffold to which the cells of a tissue are anchored

B-cells – these are cells of the adaptive immune system which produce antibodies that can bind with molecules projecting from the surfaces of non-self cells.

Beta-catenin – TLR4 activation in NEC decreases cell renewal through this pathway in isolated small intestine IECs; this protein is also a subunit of the cadherin cell junction complexes and intracellular transducer of WNT signaling pathway; Drosophila has a homolog called armadillo.

Beta-cellulin – an EFGR ligand that plays a role in gut epithelial cell proliferation, differentiation and migration

CCL20 – this cytokine is important for recruiting lymphocytes and dendritic cells into the lamina propria; it is a chemoattractant for these cells and acts as a ligand for their CCR6 receptors.

CCR6 – chemokine receptor 6; a receptor on lymphocytes and dendritic cells that triggers cell migration along a gradient of the chemokine CCL20; chemokine gradients tell immune cells where they are needed.

Commensal microbiota – bacteria and other single-celled organisms that appear to be able to live and reproduce within the microscopic ecosystem that lines our epithelium (including our skin); some are known to have a mutualistic relationship with our bodies but others may neither harm or help us (somewhat like a bird's nest on a tree branch).

COX2 – cyclooxygenase 2; one of a set of enzymes involved in synthesis of inflammatory lipid hormones known as prostaglandins; aspirin and ibuprofen are two nonsteroidal inflammation drugs (NSAIDS) that act as inhibitors of these enzymes.

CpG-ODNs – short, single stranded, synthetically produced nucleotide chains with a much higher frequency of cytosine-guanine repeats than is typically found in vertebrates; researchers use them to trigger immune responses based on studies that found these pairs to be common in

microbes, they are therefore a type of PAMP. TLR9 is the PRR most often associated with these PAMPs in the literature.

Crypts – the lowest points between the villi; this is an important zone of cell proliferation in the gut epithelium with new cells being produced at a pretty high rate in a healthy gut and migrating toward the apex of the villi to replace dying cells as though on a conveyor belt.

Cytokines – a class of small proteins and glycoproteins that serve as ligands for the receptors on a range of different types of cells in the innate and adaptive immune systems. Each of these cells has a very specific set of receptors for detecting a unique set of cytokines allowing their responses to these cytokines to be very specific as well. Depending on the unique relationship between cytokine and its “target” cells, the recipient cells may respond by producing cytokines that will activate other immune cells, directly initiate a destru(i)3(ni)36eate oth /P <</ a3(s)-2Tmits D [(t)-7(hat)

IECs – intraepithelial cells; the single layer of cells that comprise the gut epithelium; this list includes four types: absorptive enterocytes, Paneth cells, goblet cells, enteroendocrine cells.

candofat be sensD [(oul)3(s)]ouhemubod7(he)inn-6(al)3.15 T-11(hbo)11(t)-7s ps

IFN – interferon; named for their ability to interfere with viral replication, this family of cytokines can be expressed by almost all of our cells when invaded by viruses, bacteria or other pathogens as they enhance the sensitivity of neighboring cells to viruses by enhancing their expression of the MHC molecules that these cells can use to present antigens to the immune system; they also recruit Natural Killer cells and macrophages, which are both innate immune cells that can destroy and recycle the components of infected cells; macrophages can activate the adaptive immune system.

IL-6 – stimulates IECs to proliferate when an injury occurs and may inhibit apoptosis

IL-8 – a cytokine that acts as a chemokine which is especially attractive to neutrophils whose phagocytic activity is upregulated by this cytokine; this cytokine is a ligand for the receptors CXCR1 and 2.

IL-13 - this interleukin is derived from T-helper-2 (T_H2) cells and it decreases the sensitivity of IECs to LPS (a pro-inflammatory trigger from bacteria), possibly by causing these cells to decrease the quantity of TLR-4 molecules (LPS receptors) they make.

IL-33R – interleukin 33 receptor; IL33 is a member of the IR1 cytokine superfamily and its expression is upregulated by epithelial cells under inflammatory conditions and alerts the immune system to necrosis and tissue damage; the receptor is highly expressed in cells of the innate immune system and T_H2 cells; effects have been shown to be inflammatory in some tissues and anti-inflammatory in others.

IRAK – interleukin-1 receptor-associated kinase 1; this is a kinase that, when IL1 receptor is activated, activates the NFκB pathway.

JNK – c-Jun N-terminal Kinase; kinases are intracellular enzymes that transfer phosphates from high energy molecules such as ATP to a wide range of molecules with the result of changing their activity – often turning them on or off; c-Jun is the name for the substrate that this kinase phosphorylates and the phosphorylation occurs at the N-terminus or the end with the unbound amino group (the other end of a protein or peptide is the C-terminus and it has a carboxyl end that's not engaged in a peptide bond); JNK is a key component of many stress response pathways, apoptosis, T-cell differentiation, and inflammatory responses; it's a member of the MAP kinase family.

Lamina propria – a space beneath the epithelial cells of the gut where cells of the innate (e.g. dendritic cells, macrophages, monocytes, neutrophils) and adaptive (B-cells and several kinds of T-cells) immune systems localize (following inflammation signals) or lurk waiting to intercept pathogens that penetrate the epithelium or to receive instructions from other cells to either trigger an inflammatory response or prevent one

Lectin – antimicrobial products that many of our cells have the ability to make but in the gut epithelium they're mostly derived from Paneth cells; includes REG3 gamma and beta, CRP-ductin and resistin-like molecule beta

Lipoteichoic acid – a surface associated adhesion molecule released from gram positive bacteria when they are degraded

LPS – lipopolysaccharide; one of the most well known PAMPS; a molecule derived from bacteria that our cells use to detect the presence of pathogenic bacteria and which researchers commonly use in experiments when they wish to trigger an inflammatory response

MD2 – myeloid differentiation-2 protein; this molecule forms a dimer with TLR4 and is necessary for TLR4 to respond to its ligand, LPS

Mesenchymal stromal cells – these are a kind of stem cell that reside in the lamina propria; are not very lineage specific and therefore may be able to differentiate into a wider range of adult cells but in the gut epithelium they seem to secrete factors that induce proliferation of IECs; they may also modulate the inflammatory response.

MyD88 – myeloid differentiation primary-response protein 88; MyD88 is an intracellular adaptor protein that plays an important role in TLR signal transduction. Adaptor proteins don't usually have any enzymatic activity but aid in the formation of associations between other proteins in a signal transduction pathway, enabling TLR to activate the NF-κB transcription factor. In the case of TLR2 and TLR4 activation, MyD88 recruits the IRAK pathway. MyD88 knockout mice were found to be susceptible to bacterial infection but humans don't seem to be as immunocompromised when they lack functional MyD88.

Myofibrolasts – these cells are located beneath the crypts and, along with small blood vessels also located here, are considered part of the "stem cell niche"

NEC – necrotizing enterocolitis; a disease common in prematurely born infants; rapid spread and spread of pathogenic bacteria triggers inflammatory response that destroys a region of the gut

NF-κB – nuclear factor kappa-light-chain-enhancer of activated B-cells; an intracellular protein produced by almost all cells and used in a variety of different signal transduction pathways and therefore associated with a number of diseases and disorders; it's a transcription factor that plays an important role in transducing cytokine signals.

s t e r o r t o r (e r) - T

NOD – Nucleotide-binding oligomerization domain; this is an intracellular receptor that appears to have evolved to detect the bacteria that have breached the cell and are therefore no longer visible to cell surface receptors of other cells; they can be relatively specific in terms of which kinds of gene transcription pathways they trigger; responses to NOD activation include cytokine signaling (as, for instance, to recruit natural killer or NK cells) and autophagy/apoptosis; NOD2 is known to play an important role in gut homeostasis by responding to peptidoglycan released by bacteria in the gut, triggering the kinase receptor-interacting protein 2 (RIP2) which leads to activation of NFκB; activation of this system generates antimicrobial peptides (esp. alpha defensin) and mucin which help maintain a safe distance distance between the epithelium and the microbiota; Broadly, variation in NOD activation can produce a range of responses from pro-inflammatory to anti-

the crypts in the gut epithelium. In experiments in which notch was inhibited, this population of cells went into decline and some began differentiating into secretory cells with goblet or paneth characterisitcs prematurely.

PAMP – pathogen associated molecular patterns; this acronym is used to refer to any of a number of molecules that may indicate the presence of a non-self cell; these may be signaling

TOLLIP – Toll-interacting protein; this is a protein that inhibits activity of TLR2 and TLR4 through IL-1R-associated kinases (IRAKs)

TRIF – TIR(toll interleukin receptor)-domain containing adapter-inducing interferon-beta; this is an intracellular domain found in TLRs and acts an adaptor that, when TLR3 and TLR4 are triggered, together with MyD88 activates expression of specific cytokine genes. TRIF appears to provide some specificity to the TLR-MyD88(signaling domain binding receptor)-6(e s)-2(i)3

ZO1 – zonula occludens; plays an important role in crosslinking tight junction proteins and anchoring them to the actin cytoskeleton

in situ *in vivo*

- *in vitro -*
-
- *in vivo*

-
-
- *in vivo*